



Transferring principles of solid-state and Laplace NMR to the field of *in vivo* brain MRI

João P. de Almeida Martins^{1,2}, Chantal M. W. Tax³, Filip Szczepankiewicz^{4,5}, Derek K. Jones^{3,6}, Carl-Fredrik Westin^{4,5}, Daniel Topgaard^{1,2}

- 5 ¹Division of Physical Chemistry, Department of Chemistry, Lund University, Lund, Sweden
²Random Walk Imaging AB, Lund, Sweden
³Cardiff University Brain Research Imaging Centre (CUBRIC), Cardiff University, Cardiff, United Kingdom
⁴Harvard Medical School, Boston, MA, United States
⁵Radiology, Brigham and Women's Hospital, Boston, MA, United States
10 ⁶Mary MacKillop Institute for Health Research, Australian Catholic University, Melbourne, Australia.

Correspondence to: João P. de Almeida Martins (joao.martins@fkem1.lu.se)

Abstract. Magnetic resonance imaging (MRI) is the primary method for non-invasive investigations of the human brain in health, disease, and development, but yields data that are difficult to interpret whenever the millimeter-scale voxels contain multiple microscopic tissue environments with different chemical and structural properties. We propose a clinically feasible
15 MRI framework to quantify the microscopic heterogeneity of the living human brain as spatially resolved five-dimensional relaxation-diffusion distributions by augmenting a conventional diffusion-weighted imaging sequence with signal encoding principles from multidimensional solid-state nuclear magnetic resonance (NMR) spectroscopy, relaxation-diffusion correlation methods from Laplace NMR of porous media, and Monte Carlo data inversion. The high dimensionality of the distribution space allows resolution of multiple microscopic environments within each heterogeneous voxel as well as their
20 individual characterization with novel statistical measures that combine the chemical sensitivity of the relaxation rates with the link between microstructure and the anisotropic diffusivity of tissue water.

1 Introduction

The structure of the brain is shaped by both disease and normal developments on a wide range of length scales. To measure and map the cellular architecture and molecular composition of the living human brain is a challenging experimental
25 endeavor that promises far-reaching implications for both clinical diagnosis and our understanding of normal brain function. Over the last decades, Magnetic Resonance Imaging (MRI) methods have been crucial for the progress of neuroanatomical studies (Lerch et al., 2017). Most clinical MRI applications rely on detecting ¹H nuclei of water molecules to produce three-dimensional images of the human brain with a spatial resolution on the millimeter scale. Even though the attainable resolution is clearly insufficient for direct observation of individual cells, chemical and cellular features can be investigated
30 by probing their effect on magnetic resonance observables such as nuclear relaxation rates (Halle, 2006) and the translational diffusivity (Callaghan, 2011) of water. Relaxation and diffusion parameters can thus indirectly report on various microscopic



properties, including cell density (Padhani et al., 2009), orientation of nerve fibers (Basser and Pierpaoli, 1996), and presence of nutrients (Daoust et al., 2017). Current quantitative relaxation (Tofts, 2003) and diffusion (Jones, 2010) MRI observables are exquisitely sensitive to the cellular processes associated with knowledge acquisition (Zatorre et al., 2012),
35 neuropsychiatric disorders (Kubicki et al., 2007), and different tumor types (Nilsson et al., 2018a), but suffer from poor specificity and the same experimental data may support several distinct biological scenarios (Zatorre et al., 2012).

More detailed information can be obtained by taking into account that each MRI voxel comprises hundreds of thousands of cells with potentially different properties, implying that the per-voxel signal may include contributions from multiple microenvironments with distinct values of the MRI observables. To resolve the various microenvironments within a single
40 voxel remains a highly challenging problem of vital importance for the progression of quantitative MRI studies. The signals from heterogeneous materials are often approximated as integral transformations of nonparametric distributions of relaxation rates or diffusivities (Istratov and Vyvenko, 1999), which may be estimated by Laplace inversion of data acquired as a function of the relevant experimental variable (Whittall and MacKay, 1989). Within the context of human brain MRI, the components of the distributions have been assigned to water populations residing in specific tissue microenvironments such
45 as myelin (Mackay et al., 1994) and tumors (Laule et al., 2017). The power to resolve and individually characterize the different components can be boosted by combining multiple relaxation- and diffusion-encoding blocks and analyzing the data as joint probability distributions of the relevant observables (English et al., 1991). These ideas follow the principles of multidimensional nuclear magnetic resonance (NMR) spectroscopy and form the basis for multidimensional Laplace NMR which has become routine in the field of porous media (Galvosas and Callaghan, 2010; Song, 2013) and is now being
50 combined with MRI (Zhang and Blumich, 2014; Benjamini and Basser, 2017). Recently, similar relaxation-diffusion correlation protocols have been translated to *in vivo* studies using model-based rather than nonparametric data inversion (De Santis et al., 2016; Veraart et al., 2017). So far, relaxation-diffusion correlation studies have relied on the Stejskal-Tanner experiment (Stejskal and Tanner, 1965), a spin-echo diffusion-weighted sequence that has been in use for more than 50 years and where the signal is encoded for diffusion along a single axis using a pair of collinear magnetic field gradient pulses. The
55 limitations of the conventional experimental design become apparent when considering a white matter voxel comprising anisotropic domains with multiple orientations. When projected onto the measurement axis defined by the magnetic field gradients, the combination of diffusion anisotropy and orientation dispersion gives rise to a broad distribution of effective diffusivities (Topgaard and Söderman, 2002) that is challenging to retrieve with nonparametric Laplace inversion and, most importantly, impossible to differentiate from a spread of isotropic diffusivities (Mitra, 1995). Consequently, despite the fact
60 that the relaxation-diffusion correlation yields more detailed information than conventional quantitative MRI, the inherent limitations of the Stejskal-Tanner experiment prevent unambiguous discrimination between isotropic and anisotropic contributions to the diffusivity distributions as well as model-free resolution of tissue microenvironments for heterogeneous anisotropic materials such as brain tissue.

We have recently shown that data acquisition and processing schemes for correlating isotropic and anisotropic nuclear
65 interactions in multidimensional solid-state NMR spectroscopy (Schmidt-Rohr and Spiess, 1994) can be translated to



diffusion NMR (de Almeida Martins and Topgaard, 2016), relaxation-diffusion correlation NMR (de Almeida Martins and Topgaard, 2018), and diffusion MRI (Topgaard, 2019), yielding nonparametric diffusion tensor distributions (Jian et al., 2007) with resolution of multiple isotropic and anisotropic diffusion components. These “multidimensional diffusion MRI” methods (Topgaard, 2017) rely on varying both the amplitude and orientation of the magnetic field gradients within a single
70 encoding block in order to mimic the effects of sample reorientation (Frydman et al., 1992) and rotor-synchronized radio frequency pulse sequences (Gan, 1992) in multidimensional solid-state NMR to target specific aspects of the tensorial property being investigated. Here, we incorporate these ideas into a clinically feasible relaxation-diffusion correlation MRI protocol to quantify the microscopic heterogeneity of the living human brain. The suggested acquisition and analysis protocols resolve tissue heterogeneity on a five-dimensional space of transverse relaxation rates and axisymmetric diffusion
75 tensors that report on the underlying chemical composition and microscopic geometry. Nonparametric relaxation-diffusion distributions are obtained for each voxel in the three-dimensional image using Monte Carlo data inversion to deal with the non-uniqueness of the Laplace inversion and estimate the uncertainty of quantitative parameters derived from the distributions (Prange and Song, 2009). Sub-voxel tissue environments are resolved without limiting assumptions on the number or properties of the individual components and characterized with statistical measures that have intuitive relations
80 with the local microstructure.

2 Methods

2.1 Multidimensional relaxation-diffusion encoding

Figure 1A displays a pulse sequence wherein the signal $S(\tau_E, \mathbf{b})$ from a given voxel is encoded for information about the transverse relaxation rate R_2 and diffusion tensor \mathbf{D} by the experimental variables echo time τ_E and diffusion encoding tensor
85 \mathbf{b} according to (de Almeida Martins and Topgaard, 2018)

$$\frac{S(\tau_E, \mathbf{b})}{S_0} = \int_0^{+\infty} \int_{\mathbf{D} \in \text{Sym}_3^+} P(R_2, \mathbf{D}) K(\tau_E, \mathbf{b}, R_2, \mathbf{D}) d\mathbf{D} dR_2, \quad (1)$$

where $P(R_2, \mathbf{D})$ is a joint probability distribution of R_2 and \mathbf{D} , the kernel $K(\tau_E, \mathbf{b}, R_2, \mathbf{D})$ links the analysis space (R_2, \mathbf{D}) to the acquisition space (τ_E, \mathbf{b}) , S_0 denotes the signal amplitude at $(\tau_E = 0, \mathbf{b} = 0)$, and Sym_3^+ represents the mathematical space containing all 3×3 symmetric positive-definite matrices. The two sets of magnetic field gradient waveforms define an axially-symmetric b -tensor that is parameterized by its trace (b), orientation (Θ, Φ) , and normalized anisotropy (b_A) (Eriksson
90 et al., 2015), the latter controlling the influence of diffusion anisotropy on the detected signal in a manner corresponding to the effect of the angle between the main magnetic field and the rotor spinning axis in solid-state NMR (Frydman et al., 1992). While conventional diffusion encoding is limited to a single b -tensor “shape” ($b_A = 1$), we have shown that variation of b_A enables model-free separation and quantification of the isotropic and anisotropic contributions to the diffusion tensors (de Almeida Martins and Topgaard, 2016). In this work, we used the numerically optimized gradient waveforms displayed in



95 **Figure 1B** (Sjölund et al., 2015) to generate b -tensors at four distinct values of b_{Δ} . In common with conventional diffusion MRI, our method requires a minimum echo time of ~ 50 ms to accommodate diffusion encoding, causing the signal contributions from components with $R_2 > 60 \text{ s}^{-1}$ to be reduced to less than 5% of their initial amplitude. This means that the proposed protocol would require substantial signal averaging in order to quantify the fractions of fast relaxing components, thus precluding a mapping of myelin water ($R_2 \approx 70 \text{ s}^{-1}$) – one of the primary focuses of early multi-echo MRI methods
100 (Mackay et al., 1994) – within a clinically viable time.

Throughout the signal encoding process, the relaxation and diffusion of water are both affected by molecular exchange between chemically different sites and interactions with cell membranes. Averaging all these complex effects into sets of effective relaxation rates and apparent diffusion tensors, sub-voxel composition can be reported as a collection of independent tissue microenvironments, each of which characterized by a set of (R_2, \mathbf{D}) coordinates (de Almeida Martins and
105 Topgaard, 2018). Assuming axial symmetry, the various microscopic diffusion tensors are parameterized by four independent dimensions: two eigenvalues corresponding to the axial and radial diffusivities, D_{\parallel} and D_{\perp} , and the polar and azimuthal angles, θ and ϕ , describing the orientation of \mathbf{D} relative to the laboratory frame of reference. The D_{\parallel} and D_{\perp} diffusivities can be combined to define measures of isotropic diffusivity, $D_{\text{iso}} = (D_{\parallel} + 2D_{\perp})/3$, and normalized diffusion anisotropy, $D_{\Delta} = (D_{\parallel} - D_{\perp})/3D_{\text{iso}}$ (Eriksson et al., 2015), which report on the “size” and “shape” of the corresponding
110 microscopic diffusion patterns (Topgaard, 2017). Tissue microscopic heterogeneity is therefore characterized with $P(R_2, D_{\text{iso}}, D_{\Delta}, \theta, \phi)$ distributions, whose dimensions directly correspond to those of the 5D acquisition space $(\tau_E, b, b_{\Delta}, \Theta, \Phi)$:

$$\frac{S(\tau_E, b, b_{\Delta}, \Theta, \Phi)}{S_0} = \int_0^{\infty} \int_0^{\infty} \int_{-1/2}^1 \int_0^{2\pi} K(\tau_E, b, b_{\Delta}, \Theta, \Phi, R_2, D_{\text{iso}}, D_{\Delta}, \theta, \phi) \times P(R_2, D_{\text{iso}}, D_{\Delta}, \theta, \phi) d\phi \sin \theta d\theta dD_{\Delta} dD_{\text{iso}} dR_2. \quad (2)$$

The relaxation-diffusion encoding kernel is defined as

$$K(\dots) = \exp(-\tau_E R_2) \exp\left(-b D_{\text{iso}} \left[1 + 2b_{\Delta} D_{\Delta} P_2(\cos \beta)\right]\right), \quad (3)$$

where $P_2(x) = (3x^2 - 1)/2$ denotes the 2nd Legendre polynomial, and β is the arc-angle between the major symmetry axes of \mathbf{b} and \mathbf{D} , given by $\cos \beta = \cos \Theta \cos \theta + \cos(\Phi - \phi) \sin \Theta \sin \theta$. According to Eq. (3), each $(\tau_E, b, b_{\Delta}, \Theta, \Phi)$ coordinate establishes
115 correlations across the separate dimensions of the R_2 - \mathbf{D} space. Consequently, sampling various combinations of echo times and b -tensor parameters facilitates a comprehensive mapping of tissue-specific relaxation and diffusion properties.

2.2 MRI measurements

A healthy volunteer (female, 31 years) was scanned on a Siemens Magnetom Prisma 3T system equipped with a 20-channel receiver head-coil, and capable of delivering gradients of 80 mT/m at the maximum slew rate of 200 T/(m·s). The



120 measurements were approved by a local Institutional Review Board (Partners Healthcare System), and the research subject provided written informed consent prior to participation.

Experimental data were acquired using the prototype spin-echo sequence (Lasič et al., 2014) and gradient waveforms shown in **Figure 1**. All images were recorded using a repetition time of 3 s, and an echo-planar readout with a $220 \times 220 \times 66$ mm³ field of view, spatial resolution of $2 \times 2 \times 6$ mm³, and a partial Fourier factor of 6/8. The depicted waveforms give four distinct b -tensor anisotropies ($b_{\Delta} = \{-0.5, 0.0, 0.5, 1.0\}$), which were probed at varying combinations of echo-times, b -values, and b -tensor orientations. The waveforms giving $b_{\Delta} = -0.5, 0.0$, and 0.5 (see **Figure 1B**) were calculated with a numerical optimization package (Sjölund et al., 2015) (<https://github.com/jsjol/NOW>), including compensation for the effects of concomitant gradients (Szczepankiewicz et al.). This procedure yielded a pair of asymmetric gradient waveforms lasting 30.8 ms and 25.0 ms, separated by approximately 8.0 ms. Linear encoding ($b_{\Delta} = 1$) was implemented with two separate
125 gradient waveforms; a symmetric bipolar gradient waveform whose encoding blocks lasted $\tau = 25.1$ ms and were separated by 8.0 ms (see **Figure 1B**), and a pair of $\tau = 15.1$ ms single-pulsed gradients bracketing a time-period of 13.7 ms. The spectral profile of the bipolar gradient waveform was tuned to that of the asymmetric gradient waveforms in order to reduce the influence of time-dependent diffusion (Woessner, 1963; Callaghan and Stepišnik, 1996).

A total of 852 images were recorded at different combinations of $(\tau_E, b, b_{\Delta}, \Theta, \Phi)$ throughout the entire scan time of 45
135 minutes. The acquisition protocol is summarized in **Figure 2A**. Briefly, $b_{\Delta} = 1$ was acquired over 72 directions distributed over four b -values (6, 10, 16, and 40 directions at $b = 0.1, 0.7, 1.4$, and $2 \cdot 10^9$ sm⁻², respectively), both $b_{\Delta} = -0.5$, and 0.5 were collected across 64 directions spread out over four b -values (6, 10, 16, and 32 directions at, respectively, $b = 0.1, 0.7, 1.4$, and $2 \cdot 10^9$ sm⁻²), and $b_{\Delta} = 0$ was acquired for a single gradient waveform orientation, repeated 6 times over six b -values ($b = 0.1, 0.3, 0.7, 1, 1.4$, and $2 \cdot 10^9$ sm⁻²). For each (b, b_{Δ}) coordinate, the set of directions was optimized using an
140 electrostatic repulsion scheme (Bak and Nielsen, 1997; Jones et al., 1999). The various $(b, b_{\Delta}, \Theta, \Phi)$ sets were then repeatedly acquired at three different echo-times ($\tau_E = 80, 110$, and 150 ms) using the spectrally-tuned waveforms. The non-tuned Stejskal-Tanner waveform was used to acquire $b_{\Delta} = 1$ data at $\tau_E = 60$ and 80 ms. Comparison between data acquired with the bipolar and the Stejskal-Tanner gradient waveforms at $\tau_E = 80$ ms allowed us to assess the validity of the Gaussian diffusion approximation (Callaghan and Stepišnik, 1996). The acquired images were not subjected to any additional corrections (*e.g.*
145 denoising or motion correction) before data inversion.

2.3 Nonparametric Monte Carlo inversion

Algorithms designed to solve Eq. (2) have been reviewed in both general (Istratov and Vyvenko, 1999) and magnetic resonance (Mitchell et al., 2012) literature. While classical inversion methods can be successfully used to estimate the 5D $P(R_2, D_{\text{iso}}, D_{\Delta}, \theta, \phi)$ distribution, they become memory costly at the high dimensionality of our protocol. To circumvent this
150 difficulty, we introduced an inversion approach wherein our correlation space is explored through a directed iterative



algorithm, explained in ref. (de Almeida Martins and Topgaard, 2018). The algorithm starts by randomly selecting 200 points from the $(0 < \log(R_2/s^{-1}) < 1.5, -10 < \log(D_{||}/m^2s^{-1}) < -8.5, -10 < \log(D_{\perp}/m^2s^{-1}) < -8.5, 0 < \cos\theta < 1, 0 < \phi < 2\pi)$ space. A discrete $P(R_2, \mathbf{D})$ distribution is then estimated by solving Eq. (2) via a non-negative least squares algorithm. Points with non-zero weights are stored and subjected to a loop where they sequentially compete with newly generated points in the 5D distribution space. Following 20 rounds, the surviving points are selected and subjected to small random mutations. Another iteration is then initialized, wherein mutated points compete amongst themselves for 20 additional rounds in order to find the local solution with the lowest residual value. A final solution is then estimated by selecting the 10 $(R_2, D_{||}, D_{\perp}, \theta, \phi)$ coordinates with the highest weights.

The procedure described above is performed voxel-wise, resulting in an array of spatially resolved $P(R_2, D_{||}, D_{\perp}, \theta, \phi)$ discrete distributions. Owing to the stochastic nature of the inversion protocol, we may fail at retrieving a non-trivial solution, which produces a small number of randomly located black voxels in the parameter maps. To correct for this, we combine the points from each voxel with the ones from its six nearest-neighbors, subsequently fitting the set of 7×10 points to the underlying signal in order to find the 10 most likely points. The new $(R_2, D_{||}, D_{\perp}, \theta, \phi)$ set is fitted to the signal, and the resulting $P(R_2, \mathbf{D})$ is taken as the solution of the analyzed voxel. Finally, the $P(R_2, D_{||}, D_{\perp}, \theta, \phi)$ distribution is mapped onto the $(R_2, D_{iso}, D_{\Delta}, \theta, \phi)$ space.

Following the works of Prange and Song (Prange and Song, 2009), we replace traditional regularization constraints (Whittall and MacKay, 1989) with an unconstrained Monte Carlo approach that estimates voxel-wise ensembles of N distinct $P(R_2, \mathbf{D})$ solutions consistent with the primary data (de Almeida Martins and Topgaard, 2018). In this study, we estimated ensembles of $N = 96$ solutions per voxel. The level of dispersion within a given solution set characterizes the uncertainty of the inversion procedure, and can thus be used to estimate the uncertainty of any quantities derived from $P(R_2, \mathbf{D})$ (Prange and Song, 2009; de Almeida Martins and Topgaard, 2018). Readers interested in a MATLAB implementation of the algorithm are directed to our GitHub repository <https://github.com/JoaoPdAMartins/md-dmri> (Nilsson et al., 2018b).

3 Results

3.1 Spatially-resolved 5D relaxation-diffusion distributions

The proposed acquisition protocol translates into distinctive signal decay curves for each of the main components of the human brain. Indeed, voxels encompassing either white matter WM, gray matter GM, or cerebrospinal fluid CSF, are all characterized by clearly distinct signal patterns (see **Figure 2B**). The observed differences can be used to infer the gross R_2 - \mathbf{D} properties of the various cerebral constituents: WM signals are highly sensitive to both b_{Δ} and (Θ, Φ) , indicative of anisotropic diffusion along coherently aligned microscopic domains; GM signal patterns are rather insensitive to b_{Δ} and (Θ, Φ) , consistent with isotropic diffusion; and CSF data decays quickly with increasing b while remaining mostly unaffected by the other acquisition variables, features that suggest an isotropic medium characterized by relatively low R_2 -values.



Voxels comprising mixtures of WM, GM, and/or CSF generate patterns that can be interpreted as a superposition of the signal data from the pure components.

Spatially resolved 5D R_2 - \mathbf{D} nonparametric distributions are retrieved from the experimental data using the model-free inversion approach described in the Methods section. **Figure 2C** displays the solution ensembles for voxels containing WM, GM, and CSF, as well as combinations of those components: WM+GM, WM+CSF, and GM+CSF. Brain tissue possesses various microscopic components, whose relaxation and diffusion properties differ over various orders of magnitude. Therefore, tissue heterogeneity is more suitably described with logarithmic distributions, where pore anisotropy is parameterized with $\log(D_{\parallel}/D_{\perp})$ instead of D_{Δ} . The distinctive characters of the raw signal patterns in **Figure 2B** result in unique voxel-wise distributions that capture the gross microscopic features of the main cerebral components. Namely, CSF is characterized by high D_{iso} , low R_2 , and $D_{\parallel} \sim D_{\perp}$; in contrast, GM and WM both exhibit lower D_{iso} and higher R_2 , with WM being differentiated by its high D_{\parallel}/D_{\perp} . As expected, voxels comprising mixtures of WM, GM, and CSF yield a linear combination of the distributions from the individual components.

Voxels containing pure GM or WM yield bimodal and unimodal distributions, respectively, that feature clusters of points covering a significant range of the R_2 - \mathbf{D} space. Because both tissue types comprise a plethora of cells with varying geometries or chemical compositions (*e.g.* axons with various amounts of myelin, dendrites, or glial cells), the observed spread may be interpreted as a direct consequence of the underlying cellular heterogeneity. However, similar broad distributions were also observed in spectroscopic multidimensional diffusion correlation measurements of discrete-component phantoms (de Almeida Martins and Topgaard, 2016, 2018), hinting that the solution spread additionally reflects the measurement and inversion uncertainty. This intrinsic uncertainty masks the effects of finer cellular details like the intra- and extra-axonal components modeled in previous diffusion-relaxation correlation MRI methods (Veraart et al., 2017). The bimodality of the GM distribution is attributed to the fact that prolate ($D_{\Delta} > 0$, $D_{\parallel}/D_{\perp} > 1$) and oblate ($D_{\Delta} < 0$, $D_{\parallel}/D_{\perp} < 1$) diffusion tensors with similar D_{iso} yield signal patterns that are only clearly discerned when $D_{\Delta} > 0.5$ or, equivalently, $D_{\parallel}/D_{\perp} > 4$ (Eriksson et al., 2015). Earlier MRI studies demonstrated that water diffusion in GM tissue is consistent with a low, yet non-negligible, anisotropy (Assaf, 2018). Our results are consistent with those findings, with the intrinsically low anisotropy preventing a distinction between prolate or oblate solutions, and consequently producing a nearly symmetric spread of components around the $\log(D_{\parallel}/D_{\perp}) = 0$ plane.

3.2 Statistical measures of tissue heterogeneity

The R_2 - \mathbf{D} distribution ensembles provide a wealth of information that is challenging to visualize in spatially resolved datasets with large image matrices. Drawing inspiration from the field of porous media, where ensembles of distributions have been converted into ensembles of scalar parameters such as total porosity or fraction of bound fluid (Prange and Song, 2009), we extract statistical measures from the R_2 - \mathbf{D} distributions. A multitude of statistical functionals can be computed from the same distribution, meaning that the per-voxel $P(R_2, \mathbf{D})$ ensembles generate a comprehensive set of distinct voxel-



wise parameters. As shown in **Figure 3**, the Monte Carlo realizations of $P(R_2, \mathbf{D})$ are translated into ensembles of statistical
215 measures, with 96 individual estimates being extracted for each measure. For compactness, the ensembles of statistical
parameters are reduced to an average $\langle \cdot \rangle$ and a dispersion measure $\sigma[\cdot]$ that is interpreted as the uncertainty of the estimated
functional (Prange and Song, 2009). To render the results more robust to outliers, we report $\langle \cdot \rangle$ as the ensemble median and
estimate $\sigma[\cdot]$ as a median absolute deviation. The calculation of averages (as measured by the median) reduces the
underlying ensemble of solutions into a single scalar, and allows us to convey intra-voxel composition with parameter maps
220 of average mean values $\langle E[x] \rangle$, average variances $\langle \text{Var}[x] \rangle$ and average covariances $\langle \text{Cov}[x, y] \rangle$ of all the relevant dimensions
of the 5D R_2 - \mathbf{D} space (see **Figure 3**). All of the statistical measures derived in this work parameterize diffusion tensor
anisotropy with D_{Δ}^2 rather than D_{Δ} ; this is motivated by the intrinsic difficulty of distinguishing between prolate and oblate
tensors (Eriksson et al., 2015).

The three maps in the first column of **Figure 3** provide a rough spatial overview of the principal tissue types: $\langle E[R_2] \rangle$ and
225 $\langle E[D_{\text{iso}}] \rangle$ clearly identify CSF-rich areas (low $\langle E[R_2] \rangle$ and high $\langle E[D_{\text{iso}}] \rangle$), while high $\langle E[D_{\Delta}^2] \rangle$ values separate WM from the
two other main cerebral tissues. However, mean parameter maps alone cannot identify or characterize intra-voxel
heterogeneity, and their use should be complemented with dispersion measures including, but not limited to, the (co)variance
elements displayed in columns 2 and 3 of **Figure 3**. For example, voxels surrounding the ventricles do not show a truly
distinctive feature in maps of mean values but are characterized by non-zero covariance matrix elements. To understand the
origin of the non-zero values, let us focus on the WM+CSF and GM+CSF voxels indicated in **Figure 3**. The corresponding
 $P(R_2, \mathbf{D})$ distributions (displayed in **Figure 2C**) comprise two populations at distant (R_2, D_{iso}) coordinates, and both voxels are
thus characterized by high values of $\text{Var}[R_2]$ and $\text{Var}[D_{\text{iso}}]$ (see histograms of **Figure 3**). As CSF and GM are both
characterized by a low anisotropy, GM+CSF exhibits low values of $\text{Var}[D_{\Delta}^2]$; in contrast, WM+CSF displays a significant
dispersion along D_{Δ}^2 , which results in high $\text{Var}[D_{\Delta}^2]$ values. Covariance measures inform about the correlations across the
230 various dimensions of the R_2 - \mathbf{D} space. In WM+CSF distributions, for instance, higher values of diffusion anisotropy are
correlated with higher R_2 and lower D_{iso} , which results in positive $\text{Cov}[R_2, D_{\Delta}^2]$ and negative $\text{Cov}[D_{\text{iso}}, D_{\Delta}^2]$. The elevated
 $\langle \text{Var}[R_2] \rangle$ and $\langle \text{Var}[D_{\text{iso}}] \rangle$, and negative $\langle \text{Cov}[R_2, D_{\text{iso}}] \rangle$ values found in the ventricular regions are thus interpreted as a
product of sub-voxel combinations of CSF with other components. A combination of high $\langle \text{Var}[D_{\Delta}^2] \rangle$, positive
 $\langle \text{Cov}[R_2, D_{\Delta}^2] \rangle$, and negative $\langle \text{Cov}[D_{\text{iso}}, D_{\Delta}^2] \rangle$ locate WM+CSF voxels in those same regions, while low values of $\langle \text{Var}[D_{\Delta}^2] \rangle$
240 indicate the existence of deep gray matter in the vicinity of the ventricles.

The maps displayed in **Figure 3** can also be used to identify voxels containing WM+GM mixtures. Because WM and GM
distributions are characterized by similar values of R_2 and D_{iso} , WM+GM voxels result in nearly zero values of $\text{Var}[R_2]$,
 $\text{Var}[D_{\text{iso}}]$, $\text{Cov}[D_{\text{iso}}, y]$ and $\text{Cov}[R_2, y]$. Instead, WM+GM voxels are signaled by finite values of $\langle \text{Var}[D_{\Delta}^2] \rangle$, originated by the
 $\log(D_{\parallel}/D_{\perp})$ spread observed in the underlying R_2 - \mathbf{D} distribution (see the WM+GM distribution in **Figure 3C**).



245 3.3 Bin-resolved metrics of tissue heterogeneity

A more detailed picture of intra-voxel heterogeneity is obtained by dividing the distribution space into smaller subspaces ('bins'). Based on the diffusion properties of $P(R_2, \mathbf{D})$ distributions from voxels containing a single tissue type, we define three bins that loosely correspond to the main brain components (see **Figure 4A**). The 'Big' bin ($-3.5 < \log(D_{\parallel}/D_{\perp}) < 3.5$, $-8.7 < \log(D_{\text{iso}}/\text{m}^2\text{s}^{-1}) < -8$, $-0.5 < \log(R_2/\text{s}^{-1}) < 2$) contains CSF contributions, whereas the 'Thin' ($0.6 < \log(D_{\parallel}/D_{\perp}) < 3.5$, $-10 < \log(D_{\text{iso}}/\text{m}^2\text{s}^{-1}) < -8.7$, $-0.5 < \log(R_2/\text{s}^{-1}) < 2$) and 'Thick' ($-3.5 < \log(D_{\parallel}/D_{\perp}) < 0.6$, $-10 < \log(D_{\text{iso}}/\text{m}^2\text{s}^{-1}) < -8.7$, $-0.5 < \log(R_2/\text{s}^{-1}) < 2$) bins capture the signal fractions from WM and GM, respectively. The names 'Big', 'Thin', and 'Thick' are inspired by the geometric properties of the microscopic diffusion tensors that are captured by each individual bin. Visual inspection of **Figure 4B** reveals that the spatial distributions of the three bins are consistent with the expected distributions of the corresponding tissues, providing more evidence that the coarsely defined bins allow a separation of the main cerebral constituents. Parameter maps of the per-bin means of the relaxation and diffusion properties are more straightforwardly interpreted than the heterogeneity measures derived from the entire distribution space: for example, the deep gray matter inferred in the previous paragraph is easily identifiable at the center (white arrows) of the 'Thick' maps of **Figure 4B**. Further, the correlations across the various dimensions of the diffusion space allow the resolution of subtle differences in relaxation rates. Focusing on the first column of **Figure 4B**, we notice that the 'Thick' fraction exhibits a slightly lower R_2 rate than that of the 'Thin' fraction. This behavior is in accordance with previous literature (Tofts, 2003) and is consistently observed across the entire slice.

Global and bin-resolved averages for all the analyzed voxels of the entire 3D image matrix are compiled in **Figure 5**, where per-voxel average means of R_2 , D_{iso} , and D_{Δ}^2 are plotted against their respective uncertainties, $\sigma[E[R_2]]$, $\sigma[E[D_{\text{iso}}]]$, and $\sigma[E[D_{\Delta}^2]]$, and average signal amplitudes $\langle S_0 \rangle$. Although the displayed statistical analysis is restricted to mean values, similar calculations can be done using any other scalar measure derived from the 5D R_2 - \mathbf{D} distributions. Examination of the scatter plots in **Figure 5** shows that microscopic populations with low signal fractions generate statistical measures with significantly higher uncertainties. While no immediate correlation is discerned between the estimated mean values and their corresponding uncertainty, the negative correlation between uncertainty and signal fractions introduces a significant dispersion of $\langle E[x] \rangle$ at $\langle S_0 \rangle / \max(\langle S_0 \rangle) < 0.1$ (see, for example, the D_{iso} scatterplots for the "Thin" and "Thick" populations). Despite the lower precision at low $\langle S_0 \rangle$, the various average mean values are observed to be nearly constant throughout the $\langle S_0 \rangle / \max(\langle S_0 \rangle) > 0.1$ region; the only exception is $\langle E[D_{\Delta}^2] \rangle$ for the 'Thin' fraction, which shows a higher susceptibility to noise as evidenced by its positive correlation with $\langle S_0 \rangle$.

The minor differences between the relaxation rates of the 'Thin' and 'Thick' components are also observed in the scatter plots of **Figure 5**. A more detailed analysis shows that distinct R_2 -rates can be consistently detected in voxels containing GM+WM mixtures (see **Figure 6A**), where conventional 1D R_2 distributions fail to resolve the subtle differences between components (Whittall et al., 1997). The second and third columns of **Figure 6A** display mixed voxels, where the 'Thin' and



‘Thick’ populations each account for at least 30% of the total measured signal. Approximately 75% of the mixed voxels exhibit R_2 differences greater than the estimated uncertainties, thus providing evidence that the differentiation between the R_2 -rates of the two bins is indeed a meaningful result.

280 All bin-resolved $\langle E[R_2] \rangle$ plots in **Figure 5** display a secondary cluster at high R_2 -values. Inspection of **Figure 6B** reveals that the fast relaxing cluster corresponds to the non-masked extra-meningeal tissues and, for the ‘Thin’ fraction, to the pallidum (region 1 in **Figure 6B**), a major component of the basal ganglia structures located deep in the brain. We also observe that the $\langle E[R_2] \rangle$ map of the ‘Thick’ component features three main R_2 populations: high R_2 in the skull region (red voxels), low R_2 in peripheral brain regions (green voxels), and intermediate R_2 values in the inner brain regions (yellow voxels). The various
285 populations within the ‘Thick’ bin motivate the demarcation of three separate R_2 regions within the $(-3.5 < \log(D_{\parallel}/D_{\perp}) < 0.6, -10 < \log(D_{\text{iso}}/\text{m}^2\text{s}^{-1}) < -8.7)$ space; we then define the ‘Low’ $(-0.5 < \log(R_2/\text{s}^{-1}) < 1.2)$, ‘Medium’ $(1.2 < \log(R_2/\text{s}^{-1}) < 1.4)$, and ‘High’ $(1.4 < \log(R_2/\text{s}^{-1}) < 2)$ sub-bins of **Figure 6C**. To assign cerebral components to each of the sub-bins, we compare bin-resolved maps with a high-resolution longitudinal relaxation (R_1) weighted image segmented in four tissue classes: WM, cortical GM, deep GM, and CSF. As evidenced by **Figure 6C**, the spatial distributions of the ‘Low’, and ‘Medium’ sub-
290 fractions roughly correspond to the expected distributions of cortical GM, and deep GM structures, respectively. Despite the similarities between bin-resolved and segmentation maps, the former possesses a grainier appearance and seem to miss a significant portion of deep GM tissue at the center of the slice. While the grainier aspect is caused by the higher noise of the R_2 - \mathbf{D} correlation dataset, the absence of central GM is explained by the presence of anisotropic tissues in structures such as the pallidum (region 1 in **Figure 6B**) and the thalamus (region 2 in **Figure 6B**). Those two deep GM structures may then be
295 contained within the ‘Thin’ bin, and not within the ‘Thick’ bin from which we defined the R_2 sub-spaces. Joining the contributions of cortical and deep GM within a single tissue class offers further insight on the link between microscopic tissue composition and binning (see **Figure 6D**). Comparing the 3-tissue segmentation with maps of the ‘Big’, ‘Thin’, and ‘Thick’ fractions confirms that the pallidum and part of the thalamus are captured by the ‘Thin’ bin. The two maps of **Figure 6D** also demonstrate that the ‘Big’, ‘Thin’, and ‘Thick’ bins are indeed capable of resolving the contributions from CSF,
300 WM, and GM, respectively.

4 Discussion and Conclusions

The proposed framework resolves intra-voxel heterogeneity on a 5D space of transverse relaxation rates R_2 and diffusion tensor parameters $(D_{\text{iso}}, D_{\Delta}, \theta, \phi)$. Per-voxel brain composition is broken down into a non-predefined number of microscopic environments with clearly distinct relaxation and diffusion properties. The heterogeneity within a voxel is thus resolved as
305 linear combinations of independent microscopic components that can be assigned to local tissue environments; on a global scale, the sub-voxel environments can be grouped into more general tissue classes. For healthy brain tissue, the detected microenvironments were classified into three broad bins whose diffusion properties respectively match those of the main



constituents of the brain: WM, GM, and CSF. The separation between contributions from the three bins provides a clean 3D mapping of WM, GM, and CSF that agrees well with a conventional R_1 -based tissue segmentation. Unlike automatic
310 segmentation approaches, our protocol resolves tissue environments on a continuous scale defined by microscopic MRI properties, thereby allowing for a natural mapping of heterogeneity within a single tissue class, *e.g.*, resolving anisotropic and isotropic regions within the thalamus. Hence, the presented protocol shows promise for neuroanatomy studies dealing with specific microscopic features such as nerve fiber tracking or free water mapping (Pasternak et al., 2009). Within a clinical setting, the ability to disentangle different tissue signals is expected to be useful for pathological conditions
315 associated with intra-voxel tissue heterogeneity, *e.g.* tumor infiltration in surrounding brain tissue, inflammation of cerebral tissue, or replacement of myelin with free water. In the latter example, the proposed echo-times lead to an almost complete decay of the signal contributions from myelin domains, meaning that the effects of axonal demyelination would have to be probed indirectly by tracking a reduction of the signal fraction from anisotropic sub-voxel components. Besides resolving the various microscopic domains within a voxel, we were also capable of observing subtle differences in component-specific
320 relaxation rates. As mentioned before, this information is unattainable with classical R_2 distribution protocols (Whittall et al., 1997), and its extraction relies on both the high dimensionality of our diffusion-encoding scheme and the sparsity of the $(R_2, D_{\text{iso}}, D_{\Delta}, \theta, \phi)$ space (de Almeida Martins and Topgaard, 2018). The measurement of \mathbf{D} -resolved transverse relaxation rates may complement previous work on tract-specific R_1 rates (De Santis et al., 2016).

At the cellular level, the translational dynamics of water inside the human brain is shaped by interactions with
325 macromolecules and partially permeable membranes forming compartments with barrier spacings ranging from nanometers for synaptic vesicles and myelin sheaths to micrometers for the plasma membranes of the axons. The diffusion of water during the 0.1 s time-scale of MRI signal encoding is thus affected by a myriad of complex phenomena that are not explicitly accounted for in Eq. (2). Instead, we use the well-established approach of approximating the micrometer-scale water displacements as a distribution of anisotropic Gaussian contributions (Jian et al., 2007). The measured diffusivities may
330 depend on the exact choice of experimental variables if the timing parameters of the gradient waveforms match the characteristic time-scales of displacements between cellular barriers (Woessner, 1963) or molecular exchange between tissue environments with distinctly different diffusion properties (Kärger, 1969). By augmenting our acquisition protocol with an experimental dimension in which the spectral profiles of the gradient waveforms are comprehensively varied (Callaghan and Stepišnik, 1996), microscopic barrier spacings could in principle be estimated by explicitly including the effects of restricted
335 diffusion in the kernel of Eq. (2). Here we chose to minimize the influence of time-dependence by designing waveforms with similar gradient-modulation spectra.

In the previous section, we mentioned that prolate ($D_{\Delta} > 0$) and oblate ($D_{\Delta} < 0$) diffusion tensors with $|D_{\Delta}| < 0.5$ result in similar signal decays (Eriksson et al., 2015). In the absence of orientational order, diffusion tensor anisotropy is detected as a deviation from a mono-exponential signal-decay, which, to first order, is proportional to D_{Δ}^2 (Eriksson et al., 2015).
340 Consequently, the magnitude of D_{Δ} can be easily determined at moderate b -values while the sign may require data acquired



with b -values up to $4 \cdot 10^9 \text{ sm}^{-2}$ (Eriksson et al., 2015) and echo-times comparable to the ones registered in this work; currently, such acquisition parameters can only be achieved with a specialized scanner (Jones et al., 2018).

Resolving and separately characterizing intra- and extra-axonal compartments in brain tissue has been of long-standing interest in the MRI field (Does, 2018). Recently, Veraart et al. (Veraart et al., 2017) estimated subtle differences in R_2 and
345 diffusivity parameters for the intra- and extra-axonal components of human brain white matter by applying a constrained two-component model to data acquired with a conventional relaxation-diffusion correlation protocol relying on the Stejskal-Tanner experiment. The obtained R_2 -values differ with less than a factor of two while the D_{iso} -values are nearly identical and the D_{Δ} -values are 1 (by constraint) and approximately 0.5 for the intra- and extra-cellular compartments, respectively. Comparing with the non-parametric distributions in **Figure 2**, we note that components with such similar properties would
350 be virtually impossible to resolve in our minimally constrained approach despite the additional information added by the b -tensor shape dimension. The limited resolution is consistent with the fact that Eq. (2) states an ill-posed inverse problem accommodating multiple non-unique solutions – probably also including the one with two ‘Thin’ components as assumed by Veraart et al. We suggest that the unconstrained inversion could be used as a first analysis tool to define the boundaries of a more ambitious model incorporating additional information, e.g. from microanatomy studies that is not directly observable in
355 the MRI data.

This work introduces and demonstrates a novel MRI framework, in which the microscopic heterogeneity of the living human brain is characterized via 5D correlations between the transverse relaxation rate R_2 , isotropic diffusivities D_{iso} , normalized diffusion anisotropy D_{Δ} , and diffusion tensor orientation (θ, ϕ) . The correlations allow model-free estimation of per-voxel relaxation-diffusion distributions $P(R_2, \mathbf{D})$ that combine the chemical sensitivity of R_2 with the link between microstructure
360 and the diffusion metrics. The rich information content of $P(R_2, \mathbf{D})$ is reported through a set of 21 unique maps obtained by binning and parameter calculation in the 5D distribution space. Being specific to different tissue types while relying on few assumptions, the presented protocol shows promise for explorative neuroscience studies in which microscopic tissue composition cannot be presumed *a priori*. While the data in the main text is acquired with a 45 min protocol that is incompatible with the time frame of clinical MRI, the supplementary material includes parameter maps obtained with an
365 abbreviated protocol in just 15 min. By including multi-band acquisition schemes we expect to speed up the acquisition to a clinically viable time of less than 10 min (Barth et al., 2016). Furthermore, the presented framework can be merged with MRI fingerprinting methodology (Ma et al., 2013), whose pattern matching algorithms may considerably boost the data inversion speed.



370 **Data and Code availability**

The software analysis tools discussed in this paper are available for downloading from a public GitHub repository: <https://github.com/JoaoPdAMartins/md-dmri> (Nilsson et al., 2018b). The presented *in vivo* data may be directly requested from the authors.

Acknowledgments

375 The authors thank Scott Hoge for his assistance with the MRI measurements. J.P.A.M. and D.T. were financially supported by the Swedish Foundation for Strategic Research (AM13-0090, ITM17-0267) and the Swedish Research Council (2014–3910, 2018-03697). C.M.W.T. is supported by a Rubicon grant (680-50-1527) from the Netherlands Organisation for Scientific Research (NWO). F.S. and C.-F.W. are both supported by a National Institutes of Health grant (P41EB015902). D.K.J. is supported by a Wellcome Investigator Award (096646/Z/11/Z) and a Wellcome Strategic Award (104943/Z/14/Z).

380 **Author Contributions**


D.K.J., C.-F.W., and D.T. conceived the project. J.P.A.M., C.M.W.T., and F.S. designed the acquisition protocol. F.S. and C.-F.W. acquired the data. The nonparametric Monte-Carlo algorithm was designed by J.P.A.M. and D.T., and the data analysis was performed by J.P.A.M., C.M.W.T. and D.T.. J.P.A.M., and D.T. wrote the manuscript, and all authors read and reviewed the manuscript.

385 **Competing interests**

D.T. owns shares in and J.P.A.M. is partially employed by the private company Random Walk Imaging AB (Lund, Sweden), which holds patents related to the described method. All other authors declare no competing interests.



A Pulse sequence for 5D R_2 -D encoding



B Gradient waveforms for b -tensor encoding




Figure 1 Acquisition protocol for 5D relaxation-diffusion MRI. (A) Pulse sequence for acquiring images encoded for relaxation and diffusion in a 5D space defined by the echo time τ_E , and b -tensor trace b , normalized anisotropy b_Δ , and orientation (Θ, Φ) . An EPI image readout block acquires the spin-echo produced by slice-selective 90° and 180° radio-frequency pulses. The 180° pulse is encased by a pair of gradient waveforms allowing for diffusion encoding according to principles from multidimensional solid-state NMR (Topgaard, 2017) (red, green, and blue lines). The signal is encoded for the transverse relaxation rate R_2 by varying the value of τ_E . (B) Numerically optimized gradient waveforms (Sjölund et al., 2015) yielding four distinct b -tensor shapes ($b_\Delta = -0.5, 0.0, 0.5$, and 1) (Eriksson et al., 2015).




Figure 2 Representative 5D relaxation-diffusion encoded signals $S(\tau_E, b)$ and distributions $P(R_2, D)$ for selected voxels in a living human brain. (A) Acquisition scheme showing τ_E , b , b_A , Θ , and Φ as a function of acquisition point. (B) Experimental (gray circles) and fitted (black points) $S(\tau_E, b)$ signals from three representative voxels containing white matter (WM), gray matter (GM), and cerebrospinal fluid (CSF). The presented signal data was acquired according to the scheme shown in panel A and is drawn with the same horizontal axis. (C) Nonparametric R_2 - D distributions obtained for both pure (WM, GM, CSF) and mixed (WM+GM, WM+CSF, GM+CSF) voxels. The discrete distributions are reported as scatter plots in a 3D space of the logarithms of the transverse relaxation rate R_2 , isotropic diffusivity D_{iso} , and axial-radial diffusivity ratio $D_{||}/D_{\perp}$. The diffusion tensor orientation (θ, ϕ) is color-coded as $[R, G, B] = [\cos \phi \sin \theta, \sin \phi \sin \theta, \cos \theta] \cdot |D_{||} - D_{\perp}| / \max(D_{||}, D_{\perp})$ and the circle area is proportional to the statistical weight of the corresponding component. The contour lines on the sides of the plots represent projections of the 5D $P(R_2, D)$ distribution onto the respective 2D planes. Panels (B) and (C) display the signals $S(\tau_E, b)$ and corresponding $P(R_2, D)$, respectively, for the same WM, GM, and CSF voxels.




Figure 3 Statistical measures derived from the relaxation-diffusion distributions. The ensemble of 96 distinct $P(R_2, \mathbf{D})$ solutions was used to calculate means $E[x]$, variances $\text{Var}[x]$ and covariances $\text{Cov}[x, y]$ of all combinations of transverse relaxation rate R_2 , isotropic diffusivity D_{iso} , and squared anisotropy D_{Δ}^2 . The statistical measures were all derived from the entire R_2 - \mathbf{D} distribution space on a voxel-by-voxel basis. Histograms are used to represent the parameter sets calculated for three voxels containing binary mixtures of white matter WM, grey matter GM, and cerebrospinal fluid CSF. Each histogram comprises 96 estimates of a single statistical measure. The averages of statistical measures, $\langle E[x] \rangle$, $\langle \text{Var}[x] \rangle$ and $\langle \text{Cov}[x, y] \rangle$, are displayed as parameter maps whose color scales are given by the bars along the abscissas of the histograms. The crosses and arrows identify the heterogeneous voxels analyzed in the histograms; notice that the signaled points correspond to the average (as measured by the median) of the ensembles of plausible solutions shown in the histograms.




Figure 4 Parameter maps with bin-resolved means of the relaxation-diffusion distributions. (A) Division of the R_2 - \mathbf{D} distribution space into different bins. The distribution space was separated into three bins (gray volumes) named ‘Big’, ‘Thin’, and ‘Thick’ that loosely capture the diffusion features of cerebrospinal fluid CSF, white matter WM, and gray matter GM, respectively. The 3D scatter plots display the nonparametric R_2 - \mathbf{D} distributions corresponding to the CSF (top), WM (middle), and GM (bottom) voxels selected in **Figure 2**. Superquadratic tensor glyphs are used to illustrate the representative \mathbf{D} captured by each bin. (B) Parameter maps of average per-bin means (color) of transverse relaxation rate $\langle E[R_2] \rangle$, isotropic diffusivity $\langle E[D_{iso}] \rangle$, squared anisotropy $\langle E[D_A^2] \rangle$, and diffusion tensor orientation $\langle E[\text{Orientation}] \rangle$. The orientation maps (column 4) are color-coded as $[R, G, B] = [D_{xx}, D_{yy}, D_{zz}] / \max(D_{xx}, D_{yy}, D_{zz})$, where D_{ii} are the diagonal elements of laboratory-framed average diffusion tensors estimated from the various distribution bins. Brightness indicates the signal fractions corresponding to the ‘Big’ (row 1), ‘Thin’ (row 2), and ‘Thick’ (row 3) bins. The white arrows identify deep gray matter structures.






Figure 5 Uncertainty estimation of the statistical measures derived from the relaxation-diffusion distributions. 3D density (color) scatter plots show the relationship between average initial signal intensity $\langle S_0 \rangle$, the average of mean values derived from the R_2 - \mathbf{D} distributions $\langle E[x] \rangle$, and their corresponding uncertainties $\sigma[E[x]]$. For display purposes, signal intensity values were normalized to the maximum recorded $\langle S_0 \rangle$, $\max(\langle S_0 \rangle)$. The contour lines on the side planes show 2D projections of the point density function defining the distribution of data points. The average mean values of transverse relaxation rate $\langle E[R_2] \rangle$ (row 1), isotropic diffusivity $\langle E[D_{iso}] \rangle$ (row 2), and squared anisotropy $\langle E[D_A^2] \rangle$ (row 3) were computed from all voxels whose $\langle S_0 \rangle$ was greater than 5% of $\max(\langle S_0 \rangle)$. The resulting dataset comprises 55327 voxels spread throughout all slices of the acquired 3D volume. The uncertainties of $\langle E[R_2] \rangle$, $\langle E[D_{iso}] \rangle$, and $\langle E[D_A^2] \rangle$ correspond to the median absolute deviation between measures extracted from 96 independent solutions of Equation (2): $\sigma[E[R_2]]$, $\sigma[E[D_{iso}]]$, and $\sigma[E[D_A^2]]$, respectively. All displayed data was derived from both the entire R_2 - \mathbf{D} space (column 1), and the ‘Big’ (column 2), ‘Thin’ (column 3), and ‘Thick’ (column 4) bins defined in **Figure 4A**.




A Bin-specific relaxation values



B Bin-resolved fractions (brightness) and means (color)



C Thick component sub-binning & conventional R_1 -based segmentation



D R_2 -D space binning & R_1 segmentation




Figure 6 Per-bin relaxation properties and tissue composition. (A) Transverse relaxation properties specific to each of the ‘Thin’ (red) and ‘Thick’ (green) bins defined in **Figure 4A**. The color-coded composite images (top) and histograms (bottom) display the fractional populations and average mean transverse relaxation values $\langle E[R_2] \rangle$ of the two bins. The first column displays all of the ‘Thin’ and ‘Thick’ voxels, while the two other columns focus on ‘Thin’+‘Thick’ mixtures wherein the bin-specific $\langle E[R_2] \rangle$ values exhibit either significant (second column) or non-significant (third column) differences. (B) Bin-resolved signal fractions (brightness) and average per-bin means (color) of R_2 , and squared anisotropy D_{Δ}^2 . Regions 1 and 2 identify microstructural properties singled-out in the Results section. (C) Subdivision of the ‘Thick’ bin into three different R_2 sub-spaces. The contributions from different sub-bins are compared with a high-resolution R_1 -weighted image segmented into four different tissues: white matter WM, cortical gray matter GM, deep GM, and cerebrospinal fluid CSF. Additive color maps display the spatial distribution of sub-bin fractions (from low to high R_2 : green, red, blue), and of cortical (green) and deep (red) GM. (D) Color-coded composite images showing the contributions of different bins (red=Thin, green=Thick, blue=Big) and conventional R_1 -based segmentation labels (red=WM, green=cortical+deep GM, blue=CSF).



References

- Assaf, Y.: Imaging laminar structures in the gray matter with diffusion MRI, *Neuroimage*, 10.1016/j.neuroimage.2017.12.096, 2018.
- 460 Bak, M., and Nielsen, N. C.: REPULSION, a novel approach to efficient powder averaging in solid-state NMR, *J. Magn. Reson.*, 125, 132-139, 10.1006/jmre.1996.1087, 1997.
- Barth, M., Breuer, F., Koopmans, P. J., Norris, D. G., and Poser, B. A.: Simultaneous multislice (SMS) imaging techniques, *Magn. Reson. Med.*, 75, 63-81, 10.1002/mrm.25897, 2016.
- Basser, P. J., and Pierpaoli, C.: Microstructural and physiological features of tissues elucidated by quantitative-diffusion-
 465 tensor MRI, *Journal of magnetic resonance, Series B*, 111, 209-219, 10.1016/j.jmr.2011.09.022, 1996.
- Benjamini, D., and Basser, P. J.: Magnetic resonance microdynamic imaging reveals distinct tissue microenvironments, *Neuroimage*, 163, 183-196, 10.1016/j.neuroimage.2017.09.033, 2017.
- Callaghan, P. T., and Stepišnik, J.: Generalized analysis of motion using magnetic field gradients, in: *Advances in magnetic and optical resonance*, Elsevier, 325-388, 1996.
- 470 Callaghan, P. T.: *Translational dynamics & magnetic resonance*, Oxford University Press, Oxford, 2011.
- Daoust, A., Dodd, S., Nair, G., Bouraoud, N., Jacobson, S., Walbridge, S., Reich, D. S., and Koretsky, A.: Transverse relaxation of cerebrospinal fluid depends on glucose concentration, *Magn. Reson. Imaging*, 44, 72-81, 10.1016/j.mri.2017.08.001, 2017.
- de Almeida Martins, J. P., and Topgaard, D.: Two-Dimensional Correlation of Isotropic and Directional Diffusion Using
 475 NMR, *Phys. Rev. Lett.*, 116, 087601, 10.1103/PhysRevLett.116.087601, 2016.
- de Almeida Martins, J. P., and Topgaard, D.: Multidimensional correlation of nuclear relaxation rates and diffusion tensors for model-free investigations of heterogeneous anisotropic porous materials, *Sci. Rep.*, 8, 2488, 10.1038/s41598-018-19826-9, 2018.
- De Santis, S., Barazany, D., Jones, D. K., and Assaf, Y.: Resolving relaxometry and diffusion properties within the same
 480 voxel in the presence of crossing fibres by combining inversion recovery and diffusion-weighted acquisitions, *Magn. Reson. Med.*, 75, 372-380, 10.1002/mrm.25644, 2016.
- Does, M. D.: Inferring brain tissue composition and microstructure via MR relaxometry, *Neuroimage*, 182, 136-148, 10.1016/j.neuroimage.2017.12.087, 2018.
- English, A. E., Whittal, K. P., Joy, M. L. G., and Henkelman, R. M.: Quantitative two-dimensional time correlation
 485 relaxometry, *Magn. Reson. Med.*, 22, 425-434, 10.1002/mrm.1910220250, 1991.
- Eriksson, S., Lasic, S., Nilsson, M., Westin, C. F., and Topgaard, D.: NMR diffusion-encoding with axial symmetry and variable anisotropy: Distinguishing between prolate and oblate microscopic diffusion tensors with unknown orientation distribution, *J. Chem. Phys.*, 142, 104201, 10.1063/1.4913502, 2015.



- Frydman, L., Chingas, G. C., Lee, Y. K., Grandinetti, P. J., Eastman, M. A., Barrall, G. A., and Pines, A.: Variable-angle correlation spectroscopy in solid-state nuclear magnetic resonance, *J. Chem. Phys.*, 97, 4800-4808, 10.1063/1.463860, 1992.
- Galvosas, P., and Callaghan, P. T.: Multi-dimensional inverse Laplace spectroscopy in the NMR of porous media, *Comptes Rendus Physique*, 11, 172-180, 10.1016/j.crhy.2010.06.014, 2010.
- Gan, Z.: High-resolution chemical shift and chemical shift anisotropy correlation in solids using slow magic angle spinning, *J. Am. Chem. Soc.*, 114, 8307-8309, 10.1021/ja00047a062, 1992.
- 495 Halle, B.: Molecular theory of field-dependent proton spin-lattice relaxation in tissue, *Magn. Reson. Med.*, 56, 60-72, 10.1002/mrm.20919, 2006.
- Istratov, A. A., and Vyvenko, O. F.: Exponential analysis in physical phenomena, *Rev. Sci. Instrum.*, 70, 1233-1257, 10.1063/1.1149581, 1999.
- Jian, B., Vemuri, B. C., Özarslan, E., Carney, P. R., and Mareci, T. H.: A novel tensor distribution model for the diffusion-weighted MR signal, *Neuroimage*, 37, 164-176, 10.1016/j.neuroimage.2007.03.074, 2007.
- 500 Jones, D., Alexander, D., Bowtell, R., Cercignani, M., Dell'Acqua, F., McHugh, D. J., Miller, K., Palombo, M., Parker, G. J., and Rudrapatna, U. J. N.: Microstructural imaging of the human brain with a 'super-scanner': 10 key advantages of ultra-strong gradients for diffusion MRI, 182, 8-38, 10.1016/j.neuroimage.2018.05.047, 2018.
- Jones, D. K., Horsfield, M. A., and Simmons, A.: Optimal strategies for measuring diffusion in anisotropic systems by magnetic resonance imaging, *Magn. Reson. Med.*, 42, 515-525, doi:10.1002/(SICI)1522-2594(199909)42:3<515::AID-MRM14>3.0.CO;2-Q, 1999.
- 505 Jones, D. K.: Diffusion MRI, Oxford University Press, 2010.
- Kärger, J.: Zur Bestimmung der Diffusion in einem Zweibereichsystem mit Hilfe von gepulsten Feldgradienten, *Annalen der Physik*, 479, 1-4, 10.1002/andp.19694790102, 1969.
- 510 Kubicki, M., McCarley, R., Westin, C.-F., Park, H.-J., Maier, S., Kikinis, R., Jolesz, F. A., and Shenton, M. E.: A review of diffusion tensor imaging studies in schizophrenia, *J. Psychiatr. Res.*, 41, 15-30, 10.1016/j.jpsychires.2005.05.005, 2007.
- Lasič, S., Szczepankiewicz, F., Eriksson, S., Nilsson, M., and Topgaard, D.: Microanisotropy imaging: quantification of microscopic diffusion anisotropy and orientational order parameter by diffusion MRI with magic-angle spinning of the q-vector, *Frontiers in Physics*, 2, 10.3389/fphy.2014.00011, 2014.
- 515 Laule, C., Bjarnason, T. A., Vavasour, I. M., Traboulsee, A. L., Moore, G. W., Li, D. K., and MacKay, A. L.: Characterization of brain tumours with spin-spin relaxation: pilot case study reveals unique T2 distribution profiles of glioblastoma, oligodendroglioma and meningioma, *J. Neurol.*, 264, 2205-2214, 10.1007/s00415-017-8609-6, 2017.
- Lerch, J. P., van der Kouwe, A. J., Raznahan, A., Paus, T., Johansen-Berg, H., Miller, K. L., Smith, S. M., Fischl, B., and Sotiropoulos, S. N.: Studying neuroanatomy using MRI, *Nat. Neurosci.*, 20, 314-326, 10.1038/nn.4501, 2017.
- 520 Ma, D., Gulani, V., Seiberlich, N., Liu, K., Sunshine, J. L., Duerk, J. L., and Griswold, M. A.: Magnetic resonance fingerprinting, *Nature*, 495, 187-192, 10.1038/nature11971, 2013.



- Mackay, A., Whittall, K., Adler, J., Li, D., Paty, D., and Graeb, D.: In vivo visualization of myelin water in brain by magnetic resonance, *Magn. Reson. Med.*, 31, 673-677, 10.1002/mrm.1910310614, 1994.
- Mitchell, J., Chandrasekera, T. C., and Gladden, L. F.: Numerical estimation of relaxation and diffusion distributions in two
525 dimensions, *Prog. Nucl. Magn. Reson. Spectrosc.*, 62, 34-50, 10.1016/j.pnmrs.2011.07.002, 2012.
- Mitra, P. P.: Multiple wave-vector extension of the NMR pulsed-field-gradient spin-echo diffusion measurement, *Phys. Rev. B*, 51, 15074-15078, 10.1103/PhysRevB.51.15074, 1995.
- Nilsson, M., Englund, E., Szczepankiewicz, F., van Westen, D., and Sundgren, P. C.: Imaging brain tumour microstructure, *Neuroimage*, 182, 232-250, 10.1016/j.neuroimage.2018.04.075, 2018a.
- 530 Nilsson, M., Szczepankiewicz, F., Lampinen, B., Ahlgren, A., De Almeida Martins, J. P., Lasic, S., Westin, C.-F., and Topgaard, D.: An open-source framework for analysis of multidimensional diffusion MRI data implemented in MATLAB, *Proc Intl Soc Mag Reson Med*, 2018b,
- Padhani, A. R., Liu, G., Mu-Koh, D., Chenevert, T. L., Thoeny, H. C., Takahara, T., Dzik-Jurasz, A., Ross, B. D., Van Cauteren, M., Collins, D., Hammoud, D. A., Rustin, G. J. S., Taouli, B., and Choyke, P. L.: Diffusion-Weighted Magnetic
535 Resonance Imaging as a Cancer Biomarker: Consensus and Recommendations, *Neoplasia*, 11, 102-125, 10.1593/neo.81328, 2009.
- Pasternak, O., Sochen, N., Gur, Y., Intrator, N., and Assaf, Y.: Free Water Elimination and Mapping from Diffusion MRI, *Magn. Reson. Med.*, 62, 717-730, 10.1002/mrm.22055, 2009.
- Prange, M., and Song, Y. Q.: Quantifying uncertainty in NMR T2 spectra using Monte Carlo inversion, *J. Magn. Reson.*,
540 196, 54-60, 10.1016/j.jmr.2008.10.008, 2009.
- Schmidt-Rohr, K., and Spiess, H. W.: *Multidimensional solid-state NMR and polymers*, Academic Press, 1994.
- Sjölund, J., Szczepankiewicz, F., Nilsson, M., Topgaard, D., Westin, C.-F., and Knutsson, H.: Constrained optimization of gradient waveforms for generalized diffusion encoding, *J. Magn. Reson.*, 261, 157-168, 10.1016/j.jmr.2015.10.012, 2015.
- Song, Y. Q.: Magnetic resonance of porous media (MRPM): a perspective, *J. Magn. Reson.*, 229, 12-24,
545 10.1016/j.jmr.2012.11.010, 2013.
- Stejskal, E. O., and Tanner, J. E.: Spin diffusion measurements: Spin echoes in the presence of a time-dependent field gradient, *J. Chem. Phys.*, 42, 288-292, doi.org/10.1063/1.1695690, 1965.
- Szczepankiewicz, F., Westin, C. F., and Nilsson, M.: Maxwell-compensated design of asymmetric gradient waveforms for tensor-valued diffusion encoding, *Magn. Reson. Med.*, 82, 1424-1437, 10.1002/mrm.27828, 2019.
- 550 Tofts, P.: *Quantitative MRI of the Brain: Measuring Changes Caused by Disease*, John Wiley & Sons, 2003.
- Topgaard, D., and Söderman, O.: Self-diffusion in two- and three-dimensional powders of anisotropic domains: An NMR study of the diffusion of water in cellulose and starch, *J. Phys. Chem. B*, 106, 11887-11892, 10.1021/jp020130p, 2002.
- Topgaard, D.: Multidimensional diffusion MRI, *J. Magn. Reson.*, 275, 98-113, 10.1016/j.jmr.2016.12.007, 2017.
- Topgaard, D.: Diffusion tensor distribution imaging, *NMR Biomed.*, 32, e4066, 10.1002/nbm.4066, 2019.



- 555 Veraart, J., Novikov, D. S., and Fieremans, E.: TE dependent Diffusion Imaging (TEdDI) distinguishes between compartmental T2 relaxation times, *Neuroimage*, 10.1016/j.neuroimage.2017.09.030, 2017.
- Whittall, K. P., and MacKay, A. L.: Quantitative interpretation of NMR relaxation data, *J. Magn. Reson.*, 84, 134-152, 10.1016/0022-2364(89)90011-5, 1989.
- Whittall, K. P., Mackay, A. L., Graeb, D. A., Nugent, R. A., Li, D. K., and Paty, D. W.: In vivo measurement of T2
560 distributions and water contents in normal human brain, *Magn. Reson. Med.*, 37, 34-43, 10.1002/mrm.1910370107, 1997.
- Woessner, D. E.: N.M.R. spin-echo self-diffusion measurements on fluids undergoing restricted diffusion, *J. Phys. Chem.*, 67, 1365-1367, 10.1021/j100800a509, 1963.
- Zatorre, R. J., Fields, R. D., and Johansen-Berg, H.: Plasticity in gray and white: neuroimaging changes in brain structure during learning, *Nat. Neurosci.*, 15, 528-536, 10.1038/nn.3045, 2012.
- 565 Zhang, Y., and Blumich, B.: Spatially resolved D-T(2) correlation NMR of porous media, *J. Magn. Reson.*, 242, 41-48, 10.1016/j.jmr.2014.01.017, 2014.