

Exploration of the close chemical space of tryptophan and tyrosine reveals importance of hydrophobicity in CW-photo-CIDNP performances

Felix Torres¹, Alois Renn¹, Roland Riek¹

5 ¹Laboratory of Physical Chemistry, ETH Zurich, Zuerich, 8093, Switzerland

Correspondence to: Roland Riek (roland.riek@phys.chem.ethz.ch)

Abstract. Sensitivity being one of the main hurdles of Nuclear Magnetic Resonance (NMR) can be gained by polarization techniques including Chemically Induced Dynamic Nuclear Polarization (CIDNP). Kaptein demonstrated that in CIDNP the polarization arises from the formation and the recombination of a radical pair in a magnetic field. In photo-CIDNP of interest

10 here the radical pair is between a dye and the molecule to be polarized. The polarization obtained is thereby dependent on a complex interplay between the two molecules and their physicochemical properties. Here, we explore continuous wave (CW) photo-CIDNP with a set of ten tryptophan and tyrosine analogues and observe not only signal enhancement of two orders of magnitude for ¹H at 600 MHz (corresponding to 10'000 times in measurement time), but also reveal that the hydrophobicity of the molecule appears to be an important factor in the polarisation extent within CW-photo-CIDNP. Furthermore, the small

15 chemical library established indicate the existence of many photo-CIDNP active molecules.

Introduction

Despite decades of development and impressive technological improvements, sensitivity remains the main hurdle of Nuclear Magnetic Resonance (NMR) spectroscopy and imaging (Ardenkjaer-Larsen et al., 2015). Chemical induced dynamic nuclear polarization (CIDNP) enhances the sensitivity of NMR thanks to out-of-Boltzmann nuclear spin polarization. The first

20 anomalous lines related to CIDNP were serendipitously observed in 1967 independently by Ward et al. (Ward and Lawler, 1967) and Bargon et al. (Bargon et al., 1967). The radical pair mechanism was proposed by Kaptein and Oosterhoff (Kaptein and Oosterhoff, 1969) and by Closs (Closs, 1969) two years after and remains the cornerstone of the CIDNP theory ever since.

Kaptein demonstrated that the polarization arises from the formation and the recombination of a radical pair in a magnetic field. The radicals can be generated in different ways such as heating, flash photolysis, and photochemical reaction. The last

25 generation mechanism is so-called photo-CIDNP and is the one presented in this work. In photo-CIDNP, light is shined into the sample where a photosensitizer is excited and can undergo intersystem-crossing towards a triplet state. The triplet state dye reacts with a molecule of interest (M) and forms a radical pair after abstraction of an electron from that molecule. The newly formed radical pair is in a triplet state and cannot recombine due to the Pauli principle. The interplay of nuclear spin dependent electron intersystem crossing into a singlet state allowing the electron back-transfer yields to different radical pair

30 recombination kinetics depending on the nuclear spin state. Therefore, CIDNP can be used to study transient radicals that are
too short lived for EPR (Closs and Trifunac, 1969;Morozova et al., 2008;Morozova et al., 2007;Morozova et al., 2005), to
study protein structure (Kaptein et al., 1978) and folding (Hore et al., 1997;Mok et al., 2003;Mok and Hore, 2004), or to study
electron-transfer mechanism (Morozova et al., 2018;Morozova et al., 2008;Morozova et al., 2005;Morozova et al., 2003). The
radical pair mechanism is extensively described in different papers, that we recommend to the reader for a deeper understanding
35 (Goez, 1995;Morozova and Ivanov, 2019;Okuno and Cavagnero, 2017). Robert Kaptein's key role in the development of the
theory underlying the CIDNP mechanism is crystallized in the Kaptein rules which capture the theory of CIDNP into a simple
equation in order to qualitatively analyze the sign of an anomalous CIDNP line (Kaptein, 1971). According to the Kaptein's
rules, considering a radical pair composed of molecules a and b the sign of the polarization on a nucleus i belonging to a is
predicted by the following equation:

40

$$\Gamma_{ne} = \mu \varepsilon \Delta g A_i \quad (1)$$

45 Γ_{ne} is the net polarization sign of the radical a, μ and ε are Boolean values. μ is positive when the radical is formed from a
triplet precursor, and negative otherwise. ε is positive for recombination products and negative for the radical escaped or the
transfer reaction products. Δg is the sign of the g-factors difference between the two radicals, i.e. g_a-g_b , and A_i is the hyperfine
coupling constant sign of the considered nucleus i in the radical a which makes the reaction nuclear spin selective. The
Kaptein's rules equation predicts sign of polarization and reflects the complex nature of the reaction path that yields to out-of-
Boltzmann spin polarization. Therefore, it can be used for a qualitative analysis of the photo-CIDNP polarized products.
Extensive studies of the photo-CIDNP effect monitored by ultra-violet absorbing dyes such as FMN (Tsentalovich et al., 2002),
50 bipyridyl (Tsentalovich et al., 2000) or 3,3',4,4'-tetracarboxybenzophenone (TCBP) (Morozova et al., 2011), enabled a great
understanding of the photo-CIDNP theory and was applied to challenging systems. It is our long-term goal to bring CIDNP
towards a versatile and straightforward applicable tool in biomedical and biochemical research. Initially, our recent efforts
have been to push the dye absorption towards more biocompatible wavelength such as near infrared (650-900 nm). In parallel,
55 the performing of photo-CIDNP with readily handled light source is a contemporary goal (Bernarding et al., 2018). These
efforts yielded to the discovery of the Atto Thio 12 (AT12) dye which monitored continuous wave (CW)-photo-CIDNP
experiments with a promising signal-to-noise enhancement (SNE) after laser irradiation at 450 nm (Sobol et al., 2019).
Furthermore, the light source was an affordable continuous-wave laser which could be setup within a few minutes on different
Bruker spectrometers, in our case: 200 MHz Avance, 600 MHz Avance III and 700 MHz Avance Neo. On this journey to
60 establish a CW-photo-CIDNP in biomedicine, a strong dependency of the photo-CIDNP SNE on the dye-molecule couple was
observed. For instance, tryptophan is poorly polarized in the presence of AT12 but highly polarized in the presence of
fluorescein, and tyrosine is highly polarized in the presence of AT12 and less well polarized in the presence of fluorescein
(Okuno and Cavagnero, 2016;Sobol et al., 2019). These changing performances were attributed in an initial approximation to
the chemical structures of the aromatic ring and the atoms in their close vicinity yielding to different magnetic parameters, g-

value and hyperfine coupling (HFC). This assumption was corroborated by the observation of anomalous line sign alternation
65 for an oxidocyclization product of tryptophan while the dye monitoring the reaction is changed from AT12 to fluorescein
(Torres et al., 2021). This observation is related to the g-factor difference between the two molecules as we shall see, and is
thus an elegant illustration of the Kaptein's rules.

Moreover, in this work we show on the importance of side-chains in the intensity of the anomalous lines in constant wave
(CW) photo-CIDNP experiments. Although the effect of the chemical modifications on the aromatic part of the molecules are
70 well described by theory and confirmed experimentally (Kuprov et al., 2007;Kuprov and Hore, 2004), the effect of the non-
aromatic moieties are considered to be conditioning the triplet state dye quenching kinetics, as observed in Time resolved (TR)
photo-CIDNP (Saprygina et al., 2014). While the exact mechanism of polarization can be only evaluated by TR-photo-CIDNP,
in the context of biomedical NMR application aiming for the highest polarization at low micromolar molecule concentration,
CW-photo-CIDNP appears to be the method of choice suggesting the exploration of a CW-photo-CIDNP-based empirical
75 approach indicated. This work reports on this approach by screening 10 compounds with two different dyes.

Results

1.1 On the Kaptein rule of the oxidocyclization product of tryptophan α -hydroxypyrrroloindole


The distinct photo-CIDNP performances of the different dye-molecule couples was previously discussed in the literature (Sobol
80 et al., 2019;Okuno and Cavagnero, 2016). For interest here, the tryptophan presented a higher signal to noise enhancement
(SNE) when polarized upon fluorescein (chemical structure see Figure 1) irradiation when compared with the dye AT12
(chemical structure see Figure 1) as shown in Figure 1 and listed in Table 1 whereas tyrosine was better polarized in the
presence of AT12 (Figure 1) (Sobol et al., 2019). The differential effect of the dyes to trigger radical pair mechanism has been
further studied and yielded to the serendipitous observation of 3 α -hydroxypyrrroloindole (HOPI, chemical structure, see Figure
85 1) an oxidocyclization product of tryptophan which is highly polarized after irradiation in the presence of AT12 (Figure 1,
Table 1). The study of HOPI revealed surprising features such as a different polarization yields between the *cis* and *trans*
diastereoisomers, and the sign alternation of the anomalous intensities depending on the dye used to form the radical pair
(Torres et al., 2021). This sign alternation is here now assessed in the light of the Kaptein rules (eq. 1). In the case of the photo-
90 CIDNP reaction that is performed for all the experiments of this work, μ is positive since the radical pair is formed in a triplet-
state. Moreover, the polarized species are the recombination products of the radical pairs and thus the parameter ε is positive.
Hence, the variable parameters are the hyperfine coupling when the molecule changes, e.g. from tryptophan to HOPI, or/and
the Δg . The Δg can also alter when the dye used is changed. Hence, for the same molecule a sign change of the NMR signal
upon switch from a dye to another one is necessarily caused by an alternation of the sign of the Δg in the Kaptein rule equation
(eq. (1)). As shown in Figure 1, the sign switch is observed in the case of HOPI (evidently for all the resonances) when the

95 dye is altered from fluorescein to AT12. This finding can then be used to set the unknown g-factor of HOPI radicals between
the g-factors of the dyes and in respect to the known values of tryptophan and tyrosine radicals as shown in Figure 2. Moreover,
the observation of the anomalous lines signs in photo-CIDNP experiments monitored with TCBP in previous study, enabled
to rank the HOPI compounds with a g-factor between 2.0034 (fluorescein) and 2.0035 (TCBP) (Torres et al., 2021).
This is not only an elegant illustration of the Kaptein's rules, but also a witness of a g-factor evolution upon chemical
100 modification from tryptophan to its oxidocyclization product, HOPI (Figure 1). Since the g-factor originates from spin-orbit
coupling, the shape of the aromatic ring was suspected to be the main factor for an increased g-factor and improved
polarizability. This hypothesis is consistent with the results previously obtained for tyrosine which has a comparable aromatic
105 system and is preferentially polarized in the presence of AT12 (Sobol et al., 2019). Therefore, the photo-CIDNP spectrum of
2-3 dihydro-tryptophan (dH-TRP) which has the same aromatic system as HOPI (Figure 1), was recorded for both the dyes,
110 AT12 and fluorescein (Figure 1). As expected, the polarization sign switch could be observed again upon dye change,
confirming the idea that similar aromatic systems should/may yield close g-factors (as pinpointed to in Figure 1 and 2).
However, the good photo-CIDNP performance of the HOPI compound is not observed for dH-TRP as the polarization
enhancement in the presence of AT12 was only 10-fold and 17-fold in the presence of fluorescein (Table 1). This difference
115 cannot be attributed to a slight difference in the g-factor (towards the g-factor of fluorescein) because the SNE in the presence
of fluorescein did not compensate the loss in SNE in the presence of AT12 as it would be expected if the enhancement would
solely rely on a g-factor value change (Figure 1 and Table 1).

Table 1: Signal enhancement of tryptophan and tyrosine derivatives (100 μ M) after irradiation in the presence of AT12 or fluorescein (20 μ M) at 1 W for 4 seconds. The reported charges are of the diamagnetic molecules. The values for the tyrosine enhancement are taken from ref 20 and for the tryptophan and HOPI enhancement from ref 22. Nevertheless, sample conditions were identical. The log(P) is the logarithm of the partition coefficient P, and P is the ratio of the concentration of the compounds in a mixture of the two immiscible solvents octanol and water ($P=[\text{molecule}]_{\text{octanol}}/=[\text{molecule}]_{\text{water}}$).

<i>molecule</i>	<i>AT12 SNE</i>	<i>Fluorescein SNE</i>	<i>Charge</i>	<i>Log(P)</i>
<i>tryptophan</i>	18 ^a	54 ^a	0	-1.1
<i>tryptamine</i>	45	70	+1	1.6
<i>tyrosine</i>	-38 ^b	-20 ^b	0	-2.3
<i>tyramine</i>	-60	-23	+1	1.1
<i>HOPI</i>	90 ^a	-70 ^a	0	-2.1
<i>2,3-dihydro-tryptophan</i>	10	-17	0	-1.5
<i>3-(2-(piperazin)ethyl)-</i> <i>indole</i>	44	103	+1	1.9
<i>indole propionic acid</i>	45	70	-1	1.8
<i>indole acetic acid</i>	27	54	-1	1.4

^a(Torres et al., 2021); ^b(Sobol et al., 2019)



120 **Figure 1: Photo-CIDNP spectra of tryptophan or tyrosine derivatives.** The best aromatic proton's signal-to-noise enhancement (SNE) is provided and the corresponding signal is pinpointed with an arrow. All spectra were recorded after 4 sec of irradiation at 1 Watt, in the presence of either AT12 (20 μ M, irradiation wavelength 532 nm) or fluorescein (20 μ M, irradiation wavelength 450 nm), the to-be-polarized molecule is present in concentration of 100 μ M. Color code: green are the AT12 monitored SNE values and photo-CIDNP spectra, blue are the fluorescein monitored SNE values and photo-CIDNP spectra, dark are the non-irradiated 125 reference spectra. The structures are provided in their ionization state at experimental pH 7.1.




Figure 2: The sign alternation enables to locate the molecules on the g-values scale. The ranking of HOPI and dH-TRP provides fine information about their radicals g-values. For the non-elsewhere defined abbreviations: TRP stands for tryptophan, TRPA stands for tryptamine, TYR is tyrosine, TYRA is tyramine. (Figure adapted from Sobol et al. (Sobol et al., 2019))

130 **1.2 The involvement of side chain properties in the photo-CIDNP performance of tryptophan and tyrosine derivatives**


The lower performances of the dH-TRP in comparison to HOPI's despite similar g-factors turned the focus to the potential involvement of the side chain. Prior work has been done by Saprygina et al. (2014), studying the influence of N-acetylation on the quenching rate of TCBP. The replacement of the α -amine by a N-acetyl, resulted in the vanishing of the positive charge and lower quenching rates of the triplet-state photosensitizer accompanied by lower time-resolved photo-CIDNP 135 enhancements and interpreted as causative of the N-acetylation. Similarly, to this approach, side chains modifications of the same molecular species were studied in the context of CW-photo-CIDNP.

Here, first insights into the potential role of the side chain was gathered by a comparison of the tryptophan-derivative tryptamine (chemical structure see Figure 1) with tryptophan. Tryptamine differs from tryptophan by the absence of the carboxylic acid on its side chain. Indeed, improved CW-photo-CIDNP SNE is observed for tryptamine when compared with 140 tryptophan, especially after irradiation in the presence of AT12 for which a further signal enhancement of a factor of 2 is documented (Figure 1, Table 1). Since both molecules have the same aromatic system and thus similar magnetic parameters (Connor et al., 2008), and a similar reaction mechanism is expected (i.e. ET), the improved polarization of tryptamine might

be due to the charge of the molecule that differs from tryptophan by the absence of the carboxylic acid on its side chain causing potentially a change in the quenching kinetics: Fluorescein's contains a benzocarboxylate moiety of typical pKa 2-2.5 and a 145 xanthenol of pKa 6.4, (Lavis et al., 2007) and is, in the buffer of interest, twice negatively charged. AT12 is neutral in the experimental conditions (pH = 7.1), however the aromatic system is globally carrying a positive charge (Figure 1). Due to its overall positive charge, it is expected that the quenching of fluorescein by tryptamine is faster than by tryptophan, which is globally neutral in the experimental conditions. However, the strongest improvement in terms of CW-photo-CIDNP performances is for AT12 monitored experiments, despite the rather repulsive charges in play.

150 In order to elaborate further on the hypothesis of the direct potential impact of the charge of the side-chain on the SNE of AT12 monitored CW-photo-CIDNP experiments, such experiments were conducted on tyramine (Table 1), a derivative of tyrosine where the α -carboxylate moiety is absent. Tyrosine and tyramine are preferentially polarized upon irradiation in the presence of AT12 oppositely to tryptophan/triptamine, due their higher g-factor, as shown in Figure 2. The CW-photo-CIDNP SNE in the presence of AT12 is significantly higher for tyramine when compared with tyrosine (Figure 1, table 1). The minor 155 SNE enhancement for tyramine versus tyrosine photo-CIDNP experiments monitored by fluorescein could be explained by the different Δg . This experiment supports the finding that the chemical modification of side-chains can significantly improve the SNE for CW-photo-CIDNP in the presence of AT12. Next, the CW-photo-CIDNP spectra of indole propionic acid (IPA) and indole acetic acid (IAA) have been recorded. IPA is the negatively charged analogue of tryptophan (Table 1) where the α -amine is lacking. IAA is similar to IPA but the carboxylate group is closer to the aromatic ring, since it is in the β -position. 160 Unexpectedly, the IPA yielded to the same SNE as tryptamine (Table 1) whereas an interpretation of the SNE solely based on the charged, overall negative for IPA, predicted an opposite effect on the performance as compared to the positively charged tryptamine. Despite identical charge as IPA, IAA (Table 1) exhibits comparable performances as compared to tryptophan. Moreover, the 3-(2-(piperazin)ethyl)-indole (PEI) is an analogue of tryptamine where the α -amine is replaced by a piperazin moiety. In PEI, the overall charge is similar to tryptamine since the pKa of the tertiary amine is close to 4 and only the 165 secondary amine is positively charged, due to its pKa around 9. PEI yielded similar polarization performances as tryptamine upon irradiation in the presence of AT12, and showed even higher SNE for fluorescein monitored photo-CIDNP experiments. An interpretation of these results solely based on the respective overall charges therefore fail to draw any trend. Alternatively, it could be hypothesized that a different side-chains dynamic may play a role in the SNE of CW-photo-CIDNP since with the side chain alterations not only the charge of the side chain changed but also the dynamics with the tryptamine, 170 IPA, IAA, PEI, and tyramine comprise faster side chain motion than tryptophan and tyrosine. This change in dynamics is indicated by the observation that the H_β resonances are split for tryptophan and not for the tryptamine, IPA, IAA and PEI (Figure 3). The same degeneracy of the H_β chemical shifts is observed when the amine group is removed from the tyrosine to become the tyramine (Figure 3). However, this hypothesis is not supported experimentally since the dihydro-tryptophan which has a higher side-chain mobility as compared to the HOPI shows less polarization than HOPI.


175 The only summary of this first attempt to interpret the chemical space exploration is that the simultaneous presence of the α -carboxylate and the α -amine is suboptimal for CW-photo-CIDNP SNE when monitored with fluorescein or AT12 as supported by the less good polarization properties of tryptophan, tyrosine, and dH-TRP when compared with their analogues. A corollary of the presence of these α -carboxylate and α -amine is the water solubility of the small molecule and their solvation shell. This idea brings us to another difference between the different side-chains properties, which is hydrophobicity. This can be assessed
180 with the logarithm of the calculated partition coefficient between octanol and water, $\log(P)$. The evolution of the hydrophobicity within the different families of compounds and its influence on the CW-photo-CIDNP performances was therefore investigated. Within the tryptophan derivative group, i.e. tryptophan, dihydro-tryptophan, tryptamine, IAA, IPA, PEI, the increasing hydrophobicity is beneficial to the CW-photo-CIDNP performances when monitored by both fluorescein and AT12 dyes (Figure 4A and B). The same trend is suggested for the tyrosine derivatives, tyrosine and tyramine: Tyramine
185 which is more hydrophobic is better polarized, especially in the presence of AT12, than tyrosine (Table 1). The HOPI was not included in this analysis since it is rather far away from the chemical space of the two series of interest.



190 **Figure 3: H_β anomalous signals for the different molecules. Dark lines are the reference spectra and blue lines are the irradiated spectra in the presence of fluorescein, and the green lines are the irradiated spectra in the presence of AT12. Scales between the different molecules are not respected for clarity purposes.**

The positive influence of hydrophobicity on the SNE may be explained by two distinct mechanisms that also may work in concert. First, the aromatic nature of the dye-molecule interaction is favored for more hydrophobic molecules. Second, the water shell surrounding the molecule is perturbed by the different hydrophobicity of the side-chains as it can be observed from the H_β dynamics (Figure 3). Hence, the π - π stacking between the excited dye and the molecule, and therefore the orbital overlap, could be altered in a positive manner by increasing the hydrophobicity of the molecule. In summary, the hydrophobicity variation upon side-chain modification appears to have a qualitative impact on the CW-photo-CIDNP SNE unlike the charge and dynamic variation. With other words, the observed trends suggest a positive impact on the SNE for higher hydrophobicity of the molecules sharing a common aromatic moiety. While noting these findings, it must be stated (as above) that the exact nature of the polarization can only be determined by TR-photo-CIDNP. However, the presented empirical approach is regarded informative for CW-photo-CIDNP applications as the positive aspects of CW-photo-CIDNP with several
195
200

seconds of light irradiation in terms of signal to noise and easy and cheap set up is apparent. The importance on the irradiation time is indicated in Supplementary Figure 1 yielding for both the compounds tested (i.e. HOPI and TRP) an enhanced signal by a factor of 1.5 to more than 2 between one second and four second irradiation time.



205 **Figure 4: Correlation between the molecule hydrophobicity and the photo-CIDNP performances for the tryptophan derivatives. A)**
In the case where CW-photo-CIDNP is monitored by AT12. B) In the case where CW-photo-CIDNP is monitored by fluorescein. Log(P) is the logarithm of the partition coefficient, P, between octanol and water and were calculated with DataWarrior®. A statistical analysis of the trendlines using Pearson's R coefficient and the Student t-test for the hypothesis shows that for 210 AT12 (A) a t of 4.87 and a p-value of 0.008 and for fluorescein (B) a t of 2.6 and a p-value of 0.060 is obtained. Hence, in both cases, the hypothesis H0 (absence of correlation) is rejected, and the alternative hypothesis (correlation is non-null) is retained.

Conclusion

The HOPI compound offers a nice illustration of the interplay between the choice of the dye and sign of the polarization according to the Kaptein rules. The modification of the chemical structure yield to a difference in the g-factor of the molecule 215 and a relationship between this g-factor evolution and the aromatic system shape could be built without using computational methods. The compilation of CW-photo-CIDNP data for different dyes of known g-values will help to refine the g-factor scale presented in the present work. This easy ranking will help to rationalize CW-photo-CIDNP performances for new molecules of interest. Moreover, the effect of non-aromatic modifications could be observed. The side chains of the different molecules play a key role in the CW-photo-CIDNP SNE, which could not be predicted considering solely the ionic interaction monitoring 220 where the polarization performances improved gradually within the two class of derivatives, respectively. The importance of the hydrophobicity opens the way for simple chemical space exploration, since it can be simply assessed by its $\log(P)$. Once again, the presented results demonstrate on the complex interplay between molecular properties that yield to a CW-photo-CIDNP signal-to-noise enhancement. Of great interest to us, is the possibility of empirically screening the chemical space for 225 potential highly CW-photo-CIDNP-active small molecules. From this initial study, emerges the potential of making CW-

photo-CIDNP a much broader method for signal enhancement in the biomedical NMR field, and the request for the exploration of the non-aromatic chemical-space within photo-CIDNP-active molecules.

Material and Methods

230 The NMR measurements were performed at 298K either on a Bruker Avance III 600 MHz spectrometer equipped with a cryoprobe. The irradiation of AT12 samples was performed with a Coherent Verdi V10 diode pumped solid state laser emitting at a wavelength of 532 nm. The laser used for the fluorescein samples was a Thorlabs L450P1600MM, a diode laser emitting at 450 nm. The laser light was coupled (using appropriate coupling optics) into an optical fiber (Thorlabs, FG950UEC) of length 10 m and a diameter of 0.95 mm. The end of the fiber was inserted into the sample solution in a 3 mm NMR tube to a
235 depth of about 5 mm above the NMR coil region.

Tyrosine, tyramine, tryptophan, tryptamine, IPA, IAA were purchased from Sigma, dH-TRP was purchased from Akos Pharma, and PEI was purchased from ChemSpace LLC. HOPI was synthesized in-house according to the previously published protocol(Torres et al., 2021). were prepared as stock solutions of 0.2 mg/ml and 0.18 mg/ml, respectively, in a 0.1 M sodium/potassium phosphate buffer (pH 7.1) with 5% D₂O. The stock solution of AttoThio 12 (AT12) was 1 mg/ml in H₂O.

240 To prevent dye quenching the enzyme cocktail Glucose oxidase (Go, 120 kDa), catalase (Cat, 240 kDa) and D-Glucose (G, 180 Da) was used at a concentration of 14 nM for each enzyme and 2.5 mM of Glucose, as described elsewhere(Okuno and Cavagnero, 2016;Lee and Cavagnero, 2013). The stock solutions were 0.25 μ M for Go and 0.16 μ M for Cat, respectively. The glucose stock solution was 500 mM in D₂O with 0.02% NaN₃. All the samples were prepared in a 100 mM KPO₄ buffer at pH = 7.1 with either 20 μ M AT12 or 25 μ M fluorescein, and 100 μ M target molecule.

245 References

Ardenkjaer-Larsen, J. H., Boebinger, G. S., Comment, A., Duckett, S., Edison, A. S., Engelke, F., Griesinger, C., Griffin, R. G., Hilty, C., Maeda, H., Parigi, G., Prisner, T., Ravera, E., van Bentum, J., Vega, S., Webb, A., Luchinat, C., Schwalbe, H., and Frydman, L.: Facing and Overcoming Sensitivity Challenges in Biomolecular NMR Spectroscopy, *Angew Chem Int Edit*, 54, 9162-9185, 10.1002/anie.201410653, 2015.

250 Bargon, J., Fischer, H., and Johnsen, U.: Kernresonanz-Emissionslinien Wahrend Rascher Radikalreaktionen .I. Aufnahmeverfahren Und Beispiele, *Z Naturforsch Pt A*, A 22, 1551-1555, 1967.

Bernarding, J., Euchner, F., Bruns, C., Ringleb, R., Mueller, D., Trantzschel, T., Bargon, J., Bommerich, U., and Plaumann, M.: Low-cost LED-based Photo-CIDNP Enables Biocompatible Hyperpolarization of F-19 for NMR and MRI at 7T and 4.7T, *Chemphyschem*, 19, 2453-2456, 10.1002/cphc.201800570, 2018.

255 Closs, G. L.: Mechanism explaining nuclear spin polarizations in radical combination reactions, *J. Am. Chem. Soc.*, 91, 4552-4554, 1969.

260 Closs, G. L., and Trifunac, A. D.: Chemically Induced Nuclear Spin Polarization as a Tool for Determination of Spin Multiplicities of Radical-Pair Precursors, *J Am Chem Soc*, 91, 4554-4555, Doi 10.1021/Ja01044a044, 1969.

Connor, H. D., Sturgeon, B. E., Mottley, C., Sipe, H. J., and Mason, R. P.: L-tryptophan radical cation electron spin resonance studies: Connecting solution-derived hyperfine coupling constants with protein spectral interpretations, *J Am Chem Soc*, 130, 6381-6387, 10.1021/ja0780277, 2008.

265 Hore, P. J., Winder, S. L., Roberts, C. H., and Dobson, C. M.: Stopped-Flow Photo-CIDNP Observation of Protein Folding, *J Am Chem Soc*, 119, 5049-5050, 1997.

Kaptein, R., and Oosterhoff, J. L.: Chemically induced dynamic nuclear polarization II - (Relation with anomalous ESR spectra), *Chem Phys Lett*, 4, 195-197, Doi 10.1016/0009-2614(69)80098-9, 1969.

Kaptein, R.: SIMPLE RULES FOR CHEMICALLY INDUCED DYNAMIC NUCLEAR POLARIZATION, *Journal of the Chemical Society D-Chemical Communications*, 732-733, 270 10.1039/c29710000732, 1971.

Kaptein, R., Dijkstra, K., and Nicolay, K.: Laser Photo-Cidnp as a Surface Probe for Proteins in Solution, *Nature*, 274, 293-294, Doi 10.1038/274293a0, 1978.

Kuprov, I., and Hore, P. J.: Chemically amplified 19F-1H nuclear Overhauser effects, *J Magn Reson*, 168, 1-7, 10.1016/j.jmr.2004.01.011, 2004.

275 Kuprov, I., Craggs, T. D., Jackson, S. E., and Hore, P. J.: Spin relaxation effects in photochemically induced dynamic nuclear polarization spectroscopy of nuclei with strongly anisotropic hyperfine couplings, *J Am Chem Soc*, 129, 9004-9013, 10.1021/ja0705792, 2007.

Lavis, L. D., Rutkoski, T. J., and Raines, R. T.: Tuning the pK(a) of fluorescein to optimize binding assays, *Anal Chem*, 79, 6775-6782, 10.1021/ac070907g, 2007.

280 Lee, J. H., and Cavagnero, S.: A Novel Tri-Enzyme System in Combination with Laser-Driven NMR Enables Efficient Nuclear Polarization of Biomolecules in Solution, *Journal of Physical Chemistry B*, 117, 6069-6081, 10.1021/jp4010168, 2013.

Mok, K. H., Nagashima, T., Day, I. J., Jones, J. A., Jones, C. J., Dobson, C. M., and Hore, P. J.: Rapid sample-mixing technique for transient NMR and photo-CIDNP spectroscopy: applications to real-time 285 protein folding, *J Am Chem Soc*, 125, 12484-12492, 10.1021/ja036357v, 2003.

Mok, K. H., and Hore, P. J.: Photo-CIDNP NMR methods for studying protein folding, *Methods*, 34, 75-87, 10.1016/jymeth.2004.03.006, 2004.

Morozova, O. B., Yurkovskaya, A. V., Vieth, H. M., and Sagdeev, R. Z.: Intramolecular electron transfer in tryptophan-tyrosine peptide in photoinduced reaction in aqueous solution, *Journal of Physical Chemistry B*, 107, 1088-1096, 10.1021/jp0265283, 2003.

290 Morozova, O. B., Hore, P. J., Sagdeev, R. Z., and Yurkovskaya, A. V.: Intramolecular electron transfer in lysozyme studied by time-resolved chemically induced dynamic nuclear polarization, *Journal of Physical Chemistry B*, 109, 21971-21978, 10.1021/jp053394v, 2005.

Morozova, O. B., Kiryutin, A. S., Sagdeev, R. Z., and Yurkovskaya, A. V.: Electron transfer between 295 guanosine radical and amino acids in aqueous solution. 1. Reduction of guanosine radical by tyrosine, *Journal of Physical Chemistry B*, 111, 7439-7448, 10.1021/jp067722i, 2007.

Morozova, O. B., Kiryutin, A. S., and Yurkovskaya, A. V.: Electron transfer between guanosine radicals and amino acids in aqueous solution. II. Reduction of guanosine radicals by tryptophan, *Journal of Physical Chemistry B*, 112, 2747-2754, 10.1021/jp0752318, 2008.

300 Morozova, O. B., Ivanov, K. L., Kiryutin, A. S., Sagdeev, R. Z., Kochling, T., Vieth, H. M., and Yurkovskaya, A. V.: Time-resolved CIDNP: an NMR way to determine the EPR parameters of elusive radicals, *Phys Chem Chem Phys*, 13, 6619-6627, 10.1039/c0cp02449j, 2011.

305 Morozova, O. B., Panov, M. S., Fishman, N. N., and Yurkovskaya, A. V.: Electron transfer vs. proton-coupled electron transfer as the mechanism of reaction between amino acids and triplet-excited benzophenones revealed by time-resolved CIDNP, *Phys Chem Chem Phys*, 20, 21127-21135, 10.1039/c8cp03591a, 2018.

Morozova, O. B., and Ivanov, K. L.: Time-Resolved Chemically Induced Dynamic Nuclear Polarization of Biologically Important Molecules, *Chemphyschem*, 20, 197-215, 10.1002/cphc.201800566, 2019.

310 Okuno, Y., and Cavagnero, S.: Fluorescein: A Photo-CIDNP Sensitizer Enabling Hypersensitive NMR Data Collection in Liquids at Low Micromolar Concentration, *Journal of Physical Chemistry B*, 120, 715-723, 10.1021/acs.jpcb.5b12339, 2016.

Okuno, Y., and Cavagnero, S.: Photochemically Induced Dynamic Nuclear Polarization: Basic Principles and Applications, *Emagres*, 6, 283-313, 10.1002/9780470034590.emrstm1499, 2017.

315 Saprygina, N. N., Morozova, O. B., Grampp, G., and Yurkovskaya, A. V.: Effect of amino group charge on the photooxidation kinetics of aromatic amino acids, *J Phys Chem A*, 118, 339-349, 10.1021/jp4097919, 2014.

Sobol, A., Torres, F., Aicher, A., Renn, A., and Riek, R.: Atto Thio 12 as a promising dye for photo-CIDNP, *J Chem Phys*, 151, 234201, 10.1063/1.5128575, 2019.

320 Torres, F., Sobol, A., Greenwald, J., Renn, A., and Riek, R.: Molecular features toward high photo-CIDNP hyperpolarization explored through the oxidocyclization of tryptophan, *Phys Chem Chem Phys*, 2021.

Tsentalovich, Y. P., Morozova, O. B., Yurkovskaya, A. V., Hore, P. J., and Sagdeev, R. Z.: Time-resolved CIDNP and laser flash photolysis study of the photoreactions of N-acetyl histidine with 2,2'-dipyridyl in aqueous solution, *Journal of Physical Chemistry A*, 104, 6912-6916, 10.1021/jp000019o, 2000.

325 Tsentalovich, Y. P., Lopez, J. J., Hore, P. J., and Sagdeev, R. Z.: Mechanisms of reactions of flavin mononucleotide triplet with aromatic amino acids, *Spectrochim Acta A*, 58, 2043-2050, Pii S1386-1425(01)00652-7
Doi 10.1016/S1386-1425(01)00652-7, 2002.

330 Ward, H. R., and Lawler, R. G.: Nuclear Magnetic Resonance Emission and Enhanced Absorption in Rapid Organometallic Reactions, *J Am Chem Soc*, 89, 5518-5519, Doi 10.1021/Ja00997a078, 1967.