Detecting anisotropic segmental dynamics in disordered proteins by cross-correlated spin-relaxation

Clemens Kauffmann^{1,*}, Irene Ceccolini^{1,*}, Georg Kontaxis¹, and Robert Konrat¹

¹Department of Structural and Computational Biology, Max Perutz Laboratories, University of Vienna, Vienna Biocenter Campus 5, A-1030 Vienna, Austria

^{*}These authors contributed equally to this work.

Correspondence: Clemens Kauffmann (clemens.kauffmann@univie.ac.at), Robert Konrat (robert.konrat@univie.ac.at)

Abstract. Among the numerous contributions of Geoffrey Bodenhausen to NMR spectroscopy his developments in the field of spin-relaxation methodology and theory will definitely have a long lasting impact. Starting with his seminal contributions to the excitation of multiple-quantum coherences he and his group thoroughly investigated the intricate relaxation properties of these "forbidden fruits" and developed experimental techniques to reveal the relevance of previously largely ignored cross-correlated

- 5 relaxation (CCR) effects, as "the essential is invisible to the eyes". Here we consider CCR within the challenging context of intrinsically disordered proteins (IDPs) and emphasize its potential and relevance for the studies of structural dynamics of IDPs in the future years to come. Conventionally, dynamics of globularly folded proteins are modeled and understood as deviations from otherwise rigid structures tumbling in solution. However, with increasing protein flexibility, as observed for IDPs, this apparent dichotomy between structure and dynamics becomes blurred. Although complex dynamics and ensemble averaging
- 10 might impair the extraction of mechanistic details even further, spin-relaxation uniquely encodes a protein's structural memory. Due to significant methodological developments, such as high-dimensional non-uniform sampling techniques, spin-relaxation in IDPs can now be monitored in unprecedented resolution. Not embedded within a rigid globular fold, conventional ¹⁵N spin probes might not suffice to capture the inherently local nature of IDP dynamics. To better describe and understand possible segmental motions of IDPs, we propose an experimental approach to detect the signature of anisotropic segmental dynamics
- 15 by quantifying cross-correlated spin relaxation of individual ${}^{15}N^{1}H^{N}$ and ${}^{13}C'{}^{13}C^{\alpha}$ spin pairs. By adapting Geoffrey Bodenhausen's symmetrical reconversion principle to obtain zero frequency spectral density values we can define and demonstrate more sensitive means to characterize anisotropic dynamics in IDPs.

1 Introduction

Geoffrey Bodenhausen's 70th anniversary marks an ideal occasion to take a fresh look at some of his numerous contributions to spin-relaxation methodology and theory. By considering his experiments within the challenging context of intrinsically disordered proteins (IDPs), we want to emphasize their potential and relevance in the future years to come. Arguably, this rediscovery might require some collective effort, as current trends appear to point in the opposite direction. As Paul Schanda put it recently: 'the popularity of detailed spin-relaxation measurements in liquids, *en vogue* 10 or 20 years ago, is declining; [...] Even with lengthy measurements it is not easy to gain much more insight than "loops are more flexible than secondary

- structures", which often does not answer mechanistic questions.'(Schanda, 2019, p. 3-4). While intentionally exaggerated, 25 this statement does point to some of the inherent limitations of relaxation experiments. Owing to their convoluted nature, spin-relaxation reports on protein dynamics only in ambiguous terms. A variety of stochastic processes can lead to time correlation functions (TCFs) of identical shape and form(Richert and Blumen, 1994, p. 1-7). In addition, the TCF is not probed directly, only its spectral density i.e. its Fourier transform is sampled at few select frequencies. Thus, with far more detailed
- 30 structural models at hand, protein dynamics are often understood as mere perturbations of otherwise rigid bodies tumbling in solution(Lipari and Szabo, 1982; Halle and Wennerström, 1981; Clore et al., 1990; Halle, 2009). Relaxation experiments commonly employed to calculate protein structures, such as Nuclear Overhauser effects (NOEs) and paramagnetic relaxation enhancements (PREs), are usually modeled without accounting for their dynamic nature(Iwahara et al., 2004; Clore and Iwahara, 2009; Xue et al., 2009; Vögeli, 2014). In a sense, protein dynamics might appear separate from protein structure, at least 35

within the structure-function-paradigm.

However, with increasing protein flexibility this apparent dichotomy becomes blurred as structure and dynamics can no longer be considered independent of each other. While complex dynamics and ensemble averaging obfuscate mechanistic details even further, the structural information content of relaxation parameters becomes increasingly apparent. In comparison to simple population averaged quantities, such as chemical shifts or scalar couplings, spin-relaxation uniquely encodes a

- system's structural memory, i.e. the temporal persistence of concerted motions and structural arrangements. Somewhat counter-40 intuitively, spin-relaxation experiments are among the prime sources of structural information available for disordered systems. However, due to a general lack of analytical descriptions for IDP dynamics(Modig and Poulsen, 2008; Idiyatullin et al., 2001; Bussell and Eliezer, 2001; Kadeřávek et al., 2014; Khan et al., 2015), this notion has been of somewhat academic nature until the recent past. Continuous developments in molecular dynamics (MD) simulation protocols (Piana et al., 2015; Rauscher et al.,
- 45 2015; Robustelli et al., 2018; Zerze et al., 2019; Piana et al., 2020; Gopal et al., 2021; Shea et al., 2021) demonstrate how this gap can finally be bridged, allowing us to validate, refine and/or analyze dynamic ensemble representations of proteins in solution(Kämpf et al., 2018; Kümmerer et al., 2020; Salvi et al., 2016, 2017). With the necessary timescales becoming increasingly accessible(Stone et al., 2007, 2010; Salomon-Ferrer et al., 2013; Eastman et al., 2017) and the spectral resolution provided by high-dimensional NUS experiments to overcome the problem of spectral overlap(Grudziaż et al., 2018), spin-relaxation in
- 50 IDPs can be investigated in unprecedented fashion.

This aspect alone suggests a systematic reassessment and evaluation of less commonly employed experiments. Far more pressing, in our opinion, is the inherently local nature of spin-relaxation in IDPs. In contrast to folded proteins, spins in IDPs are not embedded within a fixed molecular tumbling frame. Thus, a single 15 N nucleus per residue as a dynamic probe probe might not suffice to capture the underlying motions in adequate detail. While detecting and quantifying the presence of

55 anisotropy in IDP dynamics might seem like a rather academic endeavor, it represents an important stepping stone towards the structural interpretation of other experiments. As we recently demonstrated, an appropriate estimate for the average correlation time is an important prerequisite for the angular evaluation of cross-correlated relaxation (CCR) of remote spins(Kauffmann et al., 2021).

More immediate in its structural implications would be the presence of diffusion anisotropy, which has been hypothesized

- 60 to be of substantial size even in highly disordered proteins such as α -Synuclein. Specifically, segmental tumbling of α -helical and extended chain conformations has been implied to lead to pronounced diffusion anisotropy effects for intraresidual and sequential ¹H-¹H NOEs(Ying et al., 2014; Mantsyzov et al., 2014, 2015). At the same time, the 3D GAF model(Bremi and Brüschweiler, 1997; Lienin et al., 1998) has been invoked to further rationalize the presence of anisotropic dynamics on the local scale of the peptide plane. This model has recently been reframed by Salvi et al. to analyze MD-simulated ¹⁵N
- 65 relaxation of a partially disordered protein(Salvi et al., 2017). In essence, it was demonstrated that NH^N-TCFs are welldescribed by the $C^{\alpha}C^{\alpha}$ -TCFs of the same peptide plane as long as variations of the flanking dihedral angles and NH^N-librations are accounted for. Explicit corrections for possible effects of anisotropic segmental tumbling were not required. However, noticeable deviations could be observed for the transverse ¹⁵N relaxation of the slowly moving residual α -helix. Marcellini et al. have reported pronounced diffusion anisotropy within the α -helical region of an otherwise disordered construct. Flexible
- 70 residues were affected noticeably less. It was suggested this might be due to their average orientation in the molecular tumbling frame(Marcellini et al., 2020). The SRLS model of Meirovitch, Freed et al.(Tugarinov et al., 2001; Meirovitch et al., 2006) also predicts pronounced anisotropy for α -helices and β -sheets. However, loops and terminal chain segments appear isotropic, asserting that proteins with substantial internal mobility are best represented by an isotropic global diffusion tensor(Zerbetto et al., 2011).
- Arguably, this somewhat ambiguous body of evidence illustrates the inevitable difficulties that come with extending concepts of folded proteins to IDPs. In fact, many of the above observations might very well be case-dependent. In the present study, we want to approach this question in a more agnostic manner. Are there experimental ways to better detect the signature of anisotropic dynamics in IDPs? At what level of evidence could we evoke the mental image of extended chains and α -helical segments tumbling in solution? The principal difficulty in characterizing these structural elements lies in their translational
- 80 periodicity. In an α -helix, NH^N vectors are strongly aligned along the main axis, while in an extended chain, they are oriented perpendicularly. In order to detect possible orientational biases in their relaxation behavior, additional spin probes with different orientations must be considered. While C^{α}H^{α} might be suitable for α -helices(Barnes et al., 2019), its orientation is too similar to NH^N in the extended chain conformation. Moreover, since it does not share a peptide plane with NH^N, it varies as a function of ϕ or ψ , same as the ¹H-¹H intraresidual and sequential NOEs. Spin probes within the same peptide plane and thus 85 less ambiguous orientations would certainly be preferable.

For IDPs in particular, Kadeřávek et al. have shown that the NH^N spectral density is best mapped by a combination of transversal and longitudinal CCR rates(Kadeřávek et al., 2014) employing Geoffrey Bodenhausen's symmetrical reconversion principle(Pelupessy et al., 2003, 2007). Together with Bodenhausen and coworkers, this concept was later extended to measure the zero frequency spectral density in a single experiment(Kadeřávek et al., 2015). By translating these concepts to the C'C^{α}

90 spin pair, we want to derive and demonstrate more sensitive means to detect anisotropic segmental dynamics in IDPs.

2 Theory

Our aim is to define an experimental measure for anisotropic segmental dynamics in IDPs. While the measure itself should be general, the considered source of anisotropy will be rather specific. To assess the sensitivity of the proposed protocol, we resort to the simplified image of extended chain and α -helical segments tumbling in solution as previously asserted by Manstyzov

95 et al.(Mantsyzov et al., 2014). Before considering experimental aspects, we start by defining the spectral density. Sampled at zero frequency and/or (combinations of) the involved Larmor frequencies, it constitutes the fundamental quantity of all

$$J_{\boldsymbol{u},\boldsymbol{v}}(\omega) = \int_{0}^{\infty} C_{\boldsymbol{u},\boldsymbol{v}}(t) \cos(\omega t) dt$$
(1)

with the time correlation function (TCF),

spin-relaxation experiments:

100
$$C_{\boldsymbol{u},\boldsymbol{v}}(t) = \langle P_2(\boldsymbol{u}(0) \cdot \boldsymbol{v}(t)) \rangle$$
 (2)

where $P_2(x) = 1.5x^2 - 0.5$ is the second order Legendre polynomial, u and v represent either dipolar unit vectors or principal components of chemical shift anisotropy (CSA) tensors. Note that our simplified definition of the TCF implicitly assumes that time-dependent distance fluctuations factorize and can thus be absorbed into constant coefficients. This requirement will be well-satisfied for the spins considered henceforth.

105

For most processes, the TCF can be described as a sum/distribution of exponential decays(Lipari and Szabo, 1982; Idiyatullin et al., 2001; Modig and Poulsen, 2008; Khan et al., 2015):

$$C_{\boldsymbol{u},\boldsymbol{v}}(t) = \sum_{k=0}^{N} a_k e^{-t/\tau_k}$$
(3)

Evaluating at t = 0 yields a type of normalization condition,

$$\sum_{k=0}^{N} a_k = C_{\boldsymbol{u},\boldsymbol{v}}(0) = \langle P_2(\boldsymbol{u}(0) \cdot \boldsymbol{v}(0)) \rangle$$
(4)

110 which equates to 1 for the familiar case of auto-correlation (u = v). For cross-correlation ($u \neq v$), Eq. (4) is bounded within [-0.5, 1].

The spectral density of Eq. (3) is a sum of Lorentzians

$$J_{\boldsymbol{u},\boldsymbol{v}}(\omega) = \sum_{k=0}^{N} a_k \frac{\tau_k}{1 + (\omega \tau_k)^2}$$
(5)

Note that, depending on how the TCF and the spectral density are defined, Eq. (5) might come with additional coefficients such as the familiar factor of $\frac{2}{5}$ (Lipari and Szabo, 1982). We prefer the above definitions as they highlight $J_{u,v}(\omega)$ as a weighted average. At zero frequency all τ_k are weighted equally, i.e. J(0) encodes the average correlation time. With increased frequency



Figure 1. The Lorentzian as a function of the correlation time τ and the (Larmor) frequency. The spectral density $J(\omega)$, which is modeled as a linear combination of Lorentzians, can be pictured as a weighted average of all correlation times τ . Since J(0) weights all correlation times equally, it represents the component most sensitive to correlation times > 1 ns. The magnetic field B_0 is 18.8 T (800 MHz proton Larmor frequency).

the impact of larger τ_k becomes less pronounced. This is illustrated in Fig. 1 for a selection of Larmor frequencies assuming a magnetic field strength of 18.8 T (800 MHz proton Larmor frequency).

120

Detecting anisotropy amounts to quantifying orientational biases in the a_k and τ_k . Similarly to Mantsyzov et al.(Mantsyzov et al., 2014), we will attribute these biases to relative orientations in idealized extended chain and α -helical segments. These structural elements are well-described by an axially symmetric diffusion tensor, which yields the following expression for the spectral density(Tjandra et al., 1996; Woessner, 1962):

$$J_{\boldsymbol{u},\boldsymbol{v}}(\omega) = \sum_{k=0}^{2} A_k(\boldsymbol{u},\boldsymbol{v}) \frac{\tau_k}{1 + (\omega\tau_k)^2}$$
(6)

where

125
$$a_{0} \equiv A_{0}(\boldsymbol{u}, \boldsymbol{v}) = P_{2}(\theta_{\boldsymbol{u}})P_{2}(\theta_{\boldsymbol{v}})$$

$$a_{1} \equiv A_{1}(\boldsymbol{u}, \boldsymbol{v}) = 0.75\sin(2\theta_{\boldsymbol{u}})\sin(2\theta_{\boldsymbol{v}})\cos(\phi_{\boldsymbol{u}} - \phi_{\boldsymbol{v}})$$

$$a_{2} \equiv A_{2}(\boldsymbol{u}, \boldsymbol{v}) = 0.75\sin^{2}(\theta_{\boldsymbol{u}})\sin^{2}(\theta_{\boldsymbol{v}})\cos(2\phi_{\boldsymbol{u}} - 2\phi_{\boldsymbol{v}})$$
(7)

and (θ, ϕ) denote the polar angles in the tumbling frame. The τ_k correspond to the inverted eigenvalues of the axially symmetric diffusion tensor:

130
$$\tau_k = (6D_\perp + k^2 (D_\parallel - D_\perp))^{-1} = D_\perp^{-1} (6 + k^2 (\frac{D_\parallel}{D_\perp} - 1))^{-1}$$
 (8)

with k = 0, 1, 2. With u and v alone, Eq. (6) would not allow us to distinguish between size effects in τ_k (i.e. "segment length") and orientational biases in $A_k(u, v)$ (i.e. "secondary structure"). To quantify the anisotropy $(\frac{D_{\parallel}}{D_{\perp}})$ only, we consider another interaction described by a second set of vectors (x, y) embedded with different orientations in the same tumbling frame and focus our attention on J(0) for three reasons. First and foremost, J(0) is the component most sensitive to the $\tau_k \ge 1$ ns (cf. Fig. 1) commonly associated with tumbling motions(Kämpf et al., 2018). Secondly, the zero frequency does not depend on the

135 type of nuclei involved. Lastly, J(0) allows us to define a convenient ratio,

$$\frac{J_{\boldsymbol{u},\boldsymbol{v}}(0)}{J_{\boldsymbol{x},\boldsymbol{y}}(0)} = \frac{D_{\perp}^{-1} \sum_{k=0}^{2} A_{k}(\boldsymbol{u},\boldsymbol{v})(6+k^{2}(\frac{D_{\parallel}}{D_{\perp}}-1))^{-1}}{D_{\perp}^{-1} \sum_{k=0}^{2} A_{k}(\boldsymbol{x},\boldsymbol{y})(6+k^{2}(\frac{D_{\parallel}}{D_{\perp}}-1))^{-1}} = \frac{\sum_{k=0}^{2} A_{k}(\boldsymbol{u},\boldsymbol{v})(6+k^{2}(\frac{D_{\parallel}}{D_{\perp}}-1))^{-1}}{\sum_{k=0}^{2} A_{k}(\boldsymbol{x},\boldsymbol{y})(6+k^{2}(\frac{D_{\parallel}}{D_{\perp}}-1))^{-1}}$$
(9)

such that the explicit size dependency cancels out. In general, if (u,v) and (x,y) have fixed relative orientations and experience the same dynamics, we obtain the isotropic edge case:

140
$$\frac{J_{\boldsymbol{u},\boldsymbol{v}}(0)}{J_{\boldsymbol{x},\boldsymbol{y}}(0)} = \frac{P_2(\boldsymbol{u}\cdot\boldsymbol{v})}{P_2(\boldsymbol{x}\cdot\boldsymbol{y})}$$
(10)

Thus, the ratio (9) encodes a simple and intuitive relation: Isotropic motions tend towards the limit (10), while anisotropic motions deviate from it. Importantly, the source of anisotropy is not essential. The invoked model of a tumbling symmetric top, Eq. (6), does not need to apply in any strict sense, rather it is used to assess the sensitivity of the J(0) ratio (9) to its associated features. Presuming that segmental arrangements are sufficiently stable, remnants of Eq. (6) might still be detectable even in case of pronounced local mobility.

145

150

In general, assessing the effects of internal motions on Eq. (6) is not straightforward. Even for folded proteins, the commonly employed model-free (MF)(Lipari and Szabo, 1982) and extended MF(Clore et al., 1990) approaches are not strictly applicable in case of anisotropic diffusion(Daragan and Mayo, 1999; Halle, 2009). For CCR in particular, the proposed adaptations can become quite intricate(Vögeli, 2010). Geoffrey Bodenhausen and coworkers have investigated this topic in a series of publications(Deschamps and Bodenhausen, 2001; Deschamps, 2002; Vugmeyster et al., 2004; Abergel and Bodenhausen, 2004, 2005; Nodet et al., 2008). Given the toy nature of the considered model, we will assess the presence of fast, isotropic motions as simply as possible by introducing a fourth Lorentzian:

$$J_{\boldsymbol{u},\boldsymbol{v}}(\omega) = S^2 \sum_{k=0}^{2} A_k(\boldsymbol{u},\boldsymbol{v}) \frac{\tau_k}{1 + (\omega\tau_k)^2} + (1 - S^2) P_2(\boldsymbol{u} \cdot \boldsymbol{v}) \frac{\tau_3}{1 + (\omega\tau_3)^2}$$
(11)

with $\tau_3^{-1} = \tau_{int}^{-1} + 4D_{\perp} + 2D_{\parallel}$, where τ_{int} is the average correlation time of the fast internal motion. The generalized order 155 parameter $S^2 \in [0,1]$ acts as a weight balancing the contributions of slow anisotropic tumbling and fast isotropic motions. To account for the angular relation between u and v, a_3 necessarily corresponds to $P_2(u \cdot v)$, which follows intuitively from condition (4) assuming a fixed angle between u and v.

Of course, the additional Lorentzian can be rationalized in terms of established models. As shown in the Appendix, Eq. (11) corresponds to a simplification of the MF-like extension proposed by Ghose et al. (Ghose et al., 1998), Eq. (A1), and generalized by Vögeli and Yao (Vögeli and Yao 2009; Vögeli 2010). In this model, the generalized order parameters depend on k as well, 160 i.e. the weighting between isotropic and anisotropic contributions can indeed vary between different k. Using a single order

6

parameter is exact only if the molecule is fully rigid ($S^2 = 1$) or if the dynamics are entirely isotropic ($\frac{D_{\parallel}}{D_{\perp}} = 1$ or $S^2 = 0$). In case of pronounced diffusion anisotropy and intermediate local mobility, Eq. (11) is only an approximate interpolation between these edge cases. Still, it is worth noting that the use of a single order parameter is both a common heuristic to account for the

- 165 effects of local dynamics on experimental CCR rates(Vögeli and Yao, 2009) and an established approximation for sufficiently small angles between u and v(Tjandra et al., 1996; Kroenke et al., 1998). To keep the amount of parameters manageable, possible differences in local dynamics for different k and/or between (u,v) and (x,y) are not reflected in Eq. (11). Systematic differences in local peptide plane dynamics(Chang and Tjandra, 2005; Ferrage et al., 2006; Wang et al., 2006; Salvi et al., 2017) will necessarily result in deviations from the isotropic case, Eq. (10). While local motions could be modeled in more detail to
- 170 better match the shape of the TCF using e.g. an extended MF approach(Clore et al., 1990), correlation time distributions(Hsu et al., 2018) or dynamic detectors(Smith et al., 2017, 2019), we only intend to divide J(0), i.e. the TCF's enclosed area, into contributions with and without orientational biases in an intuitive and simple manner. Attributed solely to the A_k , Eq. (7), we now consider the effect of these biases from an experimental perspective.

3 Methods

180

175 We will assume the canonical peptide plane geometry of Corey and Pauling(Corey et al., 1953) as depicted in Fig. 2 including approximate principal components of the CSA tensors for ¹⁵N and ¹³C' adapted from Geoffrey Bodenhausen and coworkers(Cisnetti et al., 2004; Loth et al., 2005).

As demonstrated by Kadeřávek et al. (Kadeřávek et al., 2014), the spectral densities of IDPs are best mapped by combining the transversal (Γ_{xy}) and the longitudinal (Γ_z) CCR rates between the ¹⁵N CSA and the NH^N dipole. Employing the expressions of Bodenhausen and coworkers(Cisnetti et al., 2004), we have

$$\Gamma_{xy}^{N/NH} = k_{N/NH} \Delta_N \left[4J_{NH,xx}(0) + 3J_{NH,xx}(\omega_N) \right]$$
(12)

$$\Gamma_z^{N/NH} = k_{N/NH} \Delta_N \left[6J_{NH,xx}(\omega_N) \right]$$

$$k_{N/NH} = \frac{2}{5} \frac{1}{24\pi} \frac{\mu_0 \hbar \gamma_n \gamma_h}{r_{NH}^3} B_0 \gamma_n$$
(13)

where μ_0 is the vacuum permeability, \hbar is the reduced Planck constant, γ is the gyromagnetic ratio, r is the distance between the nuclei, B_0 is the magnetic field strength and $\Delta_N = (\sigma_{xx} - \sigma_{yy}) = (\sigma_{xx} - \sigma_{zz})$ is the size difference of the ¹⁵N CSA principal components (in ppm). Mapping $J_{NH,xx}(0)$ amounts to the simple subtraction $\Gamma_{xy}^{N/NH} - 0.5\Gamma_z^{N/NH}$.

To complement these rates, we consider their counterparts for the 13 C' CSA and the C'C^{α} dipole.:

$$\Gamma_{xy}^{C'/C'C^{\alpha}} = k_{C'/C'C^{\alpha}} \left[(\sigma_{xx} - \sigma_{zz}) (4J_{C'C^{\alpha},xx}(0) + 3J_{C'C^{\alpha},xx}(\omega_{C})) + (\sigma_{yy} - \sigma_{zz}) (4J_{C'C^{\alpha},yy}(0) + 3J_{C'C^{\alpha},yy}(\omega_{C})) \right]$$
(14)

$$\Gamma_{z}^{C'/C'C^{\alpha}} = k_{C'/C'C^{\alpha}} \left[(\sigma_{yy} - \sigma_{zz}) 6J_{C'C^{\alpha}xx}(\omega_{C}) + (\sigma_{yy} - \sigma_{zz}) 6J_{C'C^{\alpha}yy}(\omega_{C}) \right]$$
(15)

$$190 \quad k_{C'/C'C^{\alpha}} = \frac{2}{5} \frac{1}{24\pi} \frac{\mu_{0} \hbar \gamma_{c}^{2}}{r_{C'C^{\alpha}}^{2}} B_{0} \gamma_{c}$$

with xx,yy,zz referring to the principal components of the fully anisotropic ¹³C' CSA tensor. Again, high frequency contributions can be eliminated via the linear combination $\Gamma_{xy}^{C'/C'C^{\alpha}} - 0.5\Gamma_{z}^{C'/C'C^{\alpha}}$.



Figure 2. The peptide plane as defined by Corey and Pauling(Corey et al., 1953): $C^{\alpha}-C' = 1.53$ Å, C'-O = 1.24 Å, C'-N = 1.32 Å, N- $C^{\alpha} = 1.47$ Å. C^{α} -C'-O = 121°, C^{α} -C'-N = 114°, O-C'-N = 125°, C'-N-H = 123°, C'-N-C^{\alpha} = 123°, H-N-C^{\alpha} = 114°. N-H = 1.04 Å is taken from Ottiger and Bax (Ottiger and Bax, 1998). The ¹⁵N and ¹³C' CSA principal components are adapted from Bodenhausen and coworkers(Cisnetti et al., 2004; Loth et al., 2005). ¹⁵N: $\Delta_N \approx \sigma_{xx} - \sigma_{yy} \approx \sigma_{xx} - \sigma_{zz} = 170$ ppm, $\alpha = 20^{\circ}$. ¹³C': $\sigma_{xx} = 249.4$ ppm, $\sigma_{zz} = 170$ ppm, $\alpha = 20^{\circ}$. = 87.9 ppm, α = 37°. σ_{uu} = 191.1 ppm is obtained from the average chemical shift of Ubiquitin (BMRB 17769)(Cornilescu et al., 1998)) following the suggested calibration (Cisnetti et al., 2004). The main axis z of the diffusion anisotropy tensor is assumed to lie in the peptide plane. Its orientation is encoded by $\beta_{C^{\alpha}C'}$.

Aside from the symmetrical reconversion principle of Bodenhausen and coworkers (Pelupessy et al., 2003, 2007), ${}^{13}C'/{}^{13}C''$ CSA-DD CCR can be measured either by monitoring the relaxation asymmetry of the ${}^{13}C'{}^{13}C^{\alpha}$ doublet or by means of a 'quantitative gamma' experiment in which sum and difference of the 13 C' doublet relaxation are measured independently. In contrast to previous approaches relying on two separate experiments ('reference' and 'cross')(Schwalbe et al., 2002), we determine $\Gamma_z^{C'/C'C^{\alpha}}$ by quantifying the different longitudinal relaxation in the ${}^{13}C'{}^{13}C^{\alpha}$ doublet recorded in a non-constant-time 205 ¹³C' evolution following the relaxation period. Transverse relaxation $\Gamma_{xy}^{C'/C'C^{\alpha}}$ is measured by more conventional quantifica-

While the measurement of transverse CCR is well-established, longitudinal CCR has been studied considerably less. In part this is due to subtleties of the involved relaxation pathways which involve multi-exponential cross- and cross-correlated 195 relaxation effects. Another reason lies in technical limitations as longitudinal relaxation rates are generally smaller due to their Larmor frequency dependence. Notably, this effect is far less pronounced for the smaller correlation times present in IDPs (cf. Figure 1). While ¹⁵N¹H^N relaxation is well understood and several sensitive NMR techniques have been proposed to measure $\Gamma_{zu}^{N/NH}$ and $\Gamma_{z}^{N/NH}$ (Tiandra et al., 1996; Kroenke et al., 1998; Pelupessy et al., 2003, 2007; Kadeřávek et al., 2015), ¹³C' relaxation is generally more problematic(Dayie and Wagner, 1997; Wang et al., 2006). Since we could not find any previous attempts to measure the longitudinal CCR rate $\Gamma_z^{C'/C'C^{\alpha}}$ in the literature, we see fit to assess its general feasibility. 200



Figure 3. Experimental results from the measurements of transverse (left) and longitudinal (right) ¹³C'/¹³C^{\alpha} CSA-DD CCR as described in Sect. 3. Data were obtained using Ubiquitin, a small globular protein of 76 residues. The figure shows the spectral region for the peptide plane spanning residues I61/O62. The asymmetry of the ${}^{13}C^{\alpha}$ doublet is visible in the cross-sections taken at the positions indicated by dashed lines. As expected, CCR effects are more pronounced in the case of transverse relaxation.

To obtain sufficient spectral resolution the CCR rates are measured directly from the intensity difference in a ${}^{13}C^{\alpha}$ -coupled 3D HNCO experiment; (i) in case of transversal CCR by quantification of differential line broadening of the ${}^{13}C'{}^{13}C^{\alpha}$ doublet during constant-time 13 C' evolution and (ii) for longitudinal CCR during real-time 13 C^{α} coupled 13 C' evolution preceded by a longitudinal relaxation delay T during which 13 C'/ 13 C' 2 CSA-DD CCR is active. This approach yields reliable longitudinal

CCR rates as long as the mixing time T is short compared to 13 C' T₁ relaxation. Typical data obtained for the small globular protein Ubiquitin are shown in Fig. 3 To suppress ${}^{13}C'{}^{13}C^{\alpha}$ cross-relaxation a ${}^{13}C'$ is inverted in the middle of the relaxation delay T. Additional unwanted CCR

pathways involving the ¹³C' CSA and ¹³C'¹H/¹³C'¹⁵N dipoles are suppressed by ¹H decoupling and ¹⁵N inversion. As the ²J-215 couplings to remote carbons $({}^{13}C'{}^{13}C^{\beta}$ and ${}^{13}C'{}_{i+1}{}^{13}C^{\alpha}_{i+1})$ are not resolved, CCR effects can be expected to average out for short and intermediate mixing times. While up to 20% in size of the ${}^{13}C'{}^{13}C^{\alpha}$ CCR, the $\pm {}^{13}C'{}^{13}C^{\beta}$ CCR components contribute to the same line. The CCR rates are obtained from the ${}^{13}C'{}^{13}C^{\alpha}$ doublet as $\log(\frac{I_a}{I_b})/2T$. Details of the pulse sequence and NMR parameters will be given elsewhere. Two exemplary ${}^{13}C'{}^{13}C^{\alpha}$ doublets measured for I61/Q62 in human Ubiquitin are shown 220 in Fig. 3.

With the general feasibility of the measurements demonstrated, we can now define a ratio Q analogous to Section 2, Eq. (9):

$$Q = \frac{\Gamma_{xy}^{C'/C'C^{\alpha}} - 0.5\Gamma_{z}^{C'/C'C^{\alpha}}}{\Gamma_{xy}^{N/NH} - 0.5\Gamma_{z}^{N/NH}} = \frac{4k_{C'/C'C^{\alpha}}[(\sigma_{xx} - \sigma_{zz})J_{C'C^{\alpha},xx}(0) + (\sigma_{yy} - \sigma_{zz})J_{C'C^{\alpha},yy}(0)]}{4k_{N/NH}\Delta_{N}J_{NH,xx}(0)}$$
(16)

To assess the sensitivity of Q (16), it is evaluated according to Eqs. (7), (8),(11), (12-15) with $\tau_3^{-1} = \tau_{int}^{-1} + 4D_{\perp} + 2D_{\parallel} = \tau_{int}^{-1} + 4D_{\perp} + 2D_{\perp} = \tau_{int}^{-1} + 2D_{\perp} =$ $\tau_{int}^{-1} + \tau_{eff}^{-1}$ under the following conditions: As specified in Fig. 2, all CSA tensors have fixed orientation and size. Following Mantsyzov et al. (Mantsyzov et al., 2014), the main axis z of the axially symmetric diffusion tensor is assumed to lie in the peptide plane, hence Q is a function of the polar angles θ only, see Eq. (7). Defining the C^{α}C' orientation as 0[°] reference,

210

the main axis is rotated from 0° to 180° towards the NH^N vector assuming anisotropy values $\frac{D_{\parallel}}{D_{\perp}}$ of 1.5 and 2.5, effective tumbling times $\tau_{eff} = (4D_{\perp} + 2D_{\parallel})^{-1}$ of 1 and 2.5 ns, internal correlation times τ_{int} of 100 and 500 ps and order parameters S^2 between 0 an 1. The magnetic field strength B_0 is the same for all rates and thus cancels out. The results are summarized in Fig. 4.

4 Results and Discussion

230

Experimental considerations necessarily result in compromises. The fully anisotropic ¹³C' CSA not only leads to spectral density contributions of two perpendicular components, it is also subject to considerable variations(Markwick et al., 2005; Cisnetti et al., 2004; Loth et al., 2005). One might be tempted to avoid the uncertainties and complications that come with the ¹³C' CSA by considering dipolar relaxation only. However, compared to the NH^N spin pair, the small gyromagnetic ratios and long internucleic distances of other dipoles lead to far smaller and less sensitive rates(Carlomagno et al., 2000). In addition, the J(0) components are generally neither dominant nor easily separated. The ¹³C' CSA both provides effective means of relaxation and allows for a straightforward extraction of J(0) components. With an approximate ratio (σ_{xx} – σ_{zz})/(σ_{yy} – σ_{zz}) ≈ 1.5 and the beneficial orientation of the C'C^α vector, the J_{C'C^α,xx}(ω) contribution is generally far more pronounced:
240 For 30° ≤ α ≤ 44° the TCF amplitudes would be 0.48 ≤ P₂(C'C^α · xx) ≤ 0.79, 0.02 ≥ P₂(C'C^α · yy) ≥ -0.29 based on the geometry of Fig. 2.

Fig. 4 shows the ratio Q (16) for different choices of τ_{eff} , τ_{int} , S^2 , $\frac{D_{\parallel}}{D_{\perp}}$ as a function of the diffusion tensor orientation. The main axis is assumed to lie in the peptide plane with the orientation denoted relative to $C'C^{\alpha}$ in terms of the angle $\beta_{C'C^{\alpha}}$, see Fig. 2. Comparing all panels (a)-(f) at once, it can be seen that the isotropic ($S^2 = 0$) baseline at around 0.3 is independent of

- 245 the specific correlation times τ_{eff} and τ_{int} , see Eq. (10). The same value is obtained for $\frac{D_{\parallel}}{D_{\perp}} = 1$, which is easily assessed from the convergence behavior for different anisotropy values in (c),(f) and (a),(b),(d),(e). How strongly Q reports on the asserted presence of diffusion anisotropy depends on the S^2 -mediated weight difference between the orientation dependent $A_k \tau_k$ (7) and the isotropic τ_{eff} . Both higher overall tumbling τ_{eff} and smaller isotropic motions τ_{int} yield a more sensitive Q for increasingly smaller order parameters S^2 , see panels (a),(b),(d),(e). Of course, this interpolation is only approximate and can
- 250 become more intricate if the local motions are considered in more detail. In this simplified case, the particular choice of τ_{int} and S^2 is to some extent arbitrary as $A_k \tau_k$, τ_{int} and S^2 merely weight the isotropic and anisotropic contributions to the TCF's enclosed area J(0). Still, the range $\tau_{eff} \ge 1$ ns $> \tau_{int}$ was chosen based on timescales recently reported for MD simulations of IDPs(Kämpf et al., 2018).

Besides the obvious influence of $\frac{D_{\parallel}}{D_{\perp}}$ itself, Q strongly depends on the orientation of the diffusion tensor. The orientations of the NH^N and the C^{α}C^{α} vectors are highlighted in all panels (a)-(f). In an extended chain, C^{α}C^{α} is approximately parallel to the main axis while NH^N stands perpendicular to it or vice versa for an α -helix. Both orientations correspond well to the minimum and maximum of Q. The range of Q depends on the size of the anisotropy $\frac{D_{\parallel}}{D_{\perp}}$. For a value of 2.5, as was previously asserted for α -Synuclein(Mantsyzov et al., 2014, 2015), the effect on Q can be quite substantial, panels (a),(b),(d),(e). For $\frac{D_{\parallel}}{D_{\perp}}$ = 1.5, it is far less pronounced, panels (c),(f).



Figure 4. The ratio Q, Eq. (16), as a function of the diffusion tensor orientation denoted by $\beta_{C^{\alpha}C'}$, Fig. 2. Dashed lines indicate the orientation of the NH^N and the C^{α}C^{α} vector. All rates are calculated according to Eqs. (7), (8),(11), (12-15) with $\tau_3^{-1} = \tau_{int}^{-1} + \tau_{eff}^{-1}$. Order parameters from 0 and 1 are color-coded. Panels (a)-(f) show Q for different choices of τ_{eff} , τ_{int} and $\frac{D_{\parallel}}{D_{\perp}}$. The magnetic field strength B_0 is the same for all rates.

We conclude that, if the concept of anisotropic tumbling of segmental α-helices and extended chains is reasonably applicable and sufficiently pronounced, Q would allow to detect its signature. Actual quantification of D₁/D₁ is of course obstructed by the limited validity of the asserted dynamic model. While the presence of relaxation-active tumbling motions does imply a certain degree of local rigidity, the structural heterogeneity of IDPs certainly challenges many of the simplifying assumptions made. Still, the ratio Q might give an indication of how relevant these concepts are for different protein systems, e.g. a partially folded
region(Salvi et al., 2017; Marcellini et al., 2020) or a highly disordered segment(Mantsyzov et al., 2014). While particularly sensitive to large correlation times, Q will report on all sources of anisotropy present in J(0). Differences in local mobility, CSA tensor variations, overall structural flexibility and experimental uncertainties will certainly shift and blur the ratio expected for isotropic motions. Still, if we assume a set of consecutive residues to experience shared anisotropic diffusion, the sizes of J(0) should reflect the presence of slower tumbling motions and Q should exhibit a systematic and sequence-persistent pattern.
If Q adheres closely to the isotropic expectation, Eq. (10), it would appear that peptide plane dynamics in IDPs are isotropic and well-probed by conventional ¹⁵N relaxation alone. If Q and the sizes of J(0) imply systematic and correlated deviations

from the isotropic case, Eq. (10), it would hint towards different inherent mobilities in the peptide plane such as the 3D GAFtype(Bremi and Brüschweiler, 1997) dynamics predicted by Salvi et al. (Salvi et al., 2017). Indeed, the sensitivity of Q along the $C^{\alpha}C^{\alpha}$ direction illustrates the potential of C'/C'C^{\alpha}CCR in investigating this type of motion.

- 275
- Thus, while introduced and assessed in terms of diffusion anisotropy, we expect the combination of transversal and longitudinal $C'/C^{\alpha}C'$ CCR rates to prove informative even outside this limited scope. For the locally dominated dynamics of IDPs in particular, differences and similarities to the NH^N spectral density can provide valuable structural insights even without invoking specific dynamic models. As the previously highlighted studies demonstrate, MD simulations can be expected to play a key role in rationalizing possible sources of anisotropy for different protein systems ranging from "fully disordered" (Mantsyzov 280 et al., 2014) to partially structured (Salvi et al., 2017; Marcellini et al., 2020). In addition, the spectral densities can also be evaluated directly. While the proposed experiments do not allow to map $J_{C'C^{\alpha},xx}$ and $J_{C'C^{\alpha},yy}$ individually, the contributions of different Larmor frequencies are fully separated. Graphical representations in particular can provide model-independent intuition about the timescales at play(Idiyatullin et al., 2001; Křížová et al., 2004; Kadeřávek et al., 2014). Expressions such as $J(0) - J(\omega)$ (Idivatullin et al., 2001), intended to suppress the contribution of faster timescales (cf. Fig. 1), are available as 285 well. More extensive analysis could be realized using general frameworks such as correlation time distributions(Khan et al., 2015; Hsu et al., 2018) or dynamic detectors(Smith et al., 2017, 2019).
 - 5 Conclusions

perimental means for the investigation of anisotropic segmental dynamics in IDPs. Spectral density mapping protocols based on transversal and longitudinal CCR of NH^N were translated to the C^{α}C' spin pair of the same peptide plane. By isolating 290 and comparing the zero frequency contributions we derived an intuitive experimental measure for the presence of anisotropic dynamics in IDPs. Building on the simplified image of a symmetric top, we show that pronounced anisotropic tumbling of extended chain and α -helical segments should be readily detectable. But even outside the context of this simplistic model, contributions of different frequencies can be separated and assessed similarly to spectral density mapping protocols. Interestingly, the required measurement of longitudinal C'/C $^{\alpha}$ C' CCR has not been investigated before. Hence, a simple proof of concept for 295 a possible measurement scheme was provided. To further substantiate and explore the presented concepts in an experimental setting, a systematic evaluation of different pulse sequences is currently under preparation in our lab.

On the occasion of Geoffrey Bodenhausen's 70th anniversary, we built on his extensive body of work to conceptualize ex-

- While detecting and quantifying the presence of anisotropy in IDP dynamics might seem like a humble academic endeavor, we believe it to be an important step not only towards a better understanding of this important protein family but also towards
- 300 immediate applications in biological and biomedical research as well as drug design. We thus take particular delight from the fact that Geoffrey Bodenhausen's l'art pour l'art pulse sequence design is also a telling testimony for the unforeseeable impact of non-utilitarian basic research driven and inspired by scholarly thinking.

Appendix A

Following Ghose et al. (Ghose et al., 1998, Eq. 3, p. 488), the spectral density of an anisotropically tumbling dynamic molecule 305 can be approximated as:

$$J_{\boldsymbol{u},\boldsymbol{v}}(\omega) = \sum_{k=0}^{2} \left[\frac{S_k^2 \tau_k}{1 + (\omega \tau_k)^2} + \frac{(A_k(\boldsymbol{u}, \boldsymbol{v}) - S_k^2) \tau_3}{1 + (\omega \tau_3)^2} \right]$$
(A1)

Following the notation of (Vögeli and Yao, 2009), the order parameter is defined as

$$S_k^2 \equiv \langle A_k(\boldsymbol{u}, \boldsymbol{v}) \rangle,$$
 (A2)

i.e. as the value A_k decays to due to fast local motions. As noted by Vögeli and Yao(Vögeli and Yao, 2009), Eq. (A1) can 310 be written in the form of an auto-correlated TCF by setting A_k as a prefactor:

$$J_{\boldsymbol{u},\boldsymbol{v}}(\omega) = \sum_{k=0}^{2} A_{k}(\boldsymbol{u},\boldsymbol{v}) \left[\frac{S_{k}^{\prime 2} \tau_{k}}{1 + (\omega \tau_{k})^{2}} + \frac{(1 - S_{k}^{\prime 2}) \tau_{3}}{1 + (\omega \tau_{3})^{2}} \right]$$
(A3)

The order parameters are then defined as ratios:

$$S_k^{\prime 2} \equiv \frac{\langle A_k(\boldsymbol{u}, \boldsymbol{v}) \rangle}{A_k(\boldsymbol{u}, \boldsymbol{v})} \tag{A4}$$

While these generalized order parameters do not adhere to the definitions of Lipari and Szabo(Lipari and Szabo, 1982), they

still equate to 1 in case of a fully rigid molecule. It can be seen from Eq. (A3) that the inclusion of local motions requires one 315

order parameter for each
$$A_k$$
 in order to balance the isotropic and anisotropic contributions. The simplified spectral density
proposed in the main text, Eq. (11), corresponds to the approximation that all A_k decay about the same relative amount (as in
the edge case of isotropic tumbling). Without the index k for the order parameter, $S^2 = S_0'^2 = S_1'^2 = S_2'^2$, Eq. (A3) simplifies
to Eq. (11):

320
$$J_{\boldsymbol{u},\boldsymbol{v}}(\omega) = \sum_{k=0}^{2} A_{k}(\boldsymbol{u},\boldsymbol{v}) \left[\frac{S^{2}\tau_{k}}{1+(\omega\tau_{k})^{2}} + \frac{(1-S^{2})\tau_{3}}{1+(\omega\tau_{3})^{2}} \right] = S^{2} \sum_{k=0}^{2} A_{k}(\boldsymbol{u},\boldsymbol{v}) \frac{\tau_{k}}{1+(\omega\tau_{k})^{2}} + (1-S^{2})P_{2}(\boldsymbol{u}\cdot\boldsymbol{v}) \frac{\tau_{3}}{1+(\omega\tau_{3})^{2}}$$
(A5)

Eq. (A5) is similar to the approximation proposed by Ghose et al., (Ghose et al., 1998, Eq. 7, p. 488) but includes the prefactor $P_2(\boldsymbol{u} \cdot \boldsymbol{v})$ for the internal TCF, preserving the isotropic limit $(\frac{D_{\parallel}}{D_{\perp}} = 1, \tau_{iso} = \tau_0 = \tau_1 = \tau_2)$ as:

$$J_{\boldsymbol{u},\boldsymbol{v}}(\omega) = P_2(\boldsymbol{u}\cdot\boldsymbol{v}) \left[\frac{S^2 \tau_{iso}}{1 + (\omega\tau_{iso})^2} + \frac{(1-S^2)\tau_3}{1 + (\omega\tau_3)^2} \right]$$
(A6)

325

In this case, S^2 again corresponds to the familiar auto-correlated order parameter of Lipari and Szabo, Lipari and Szabo, 1982), the auto- and cross-correlated TCF are related by the simple factor $P_2(\boldsymbol{u} \cdot \boldsymbol{v})$. For sufficiently small angles between \boldsymbol{u} and v, this relation can be extended to approximate anisotropic tumbling in accordance with the MF rationale of Lipari and

Szabo(Tjandra et al., 1996; Kroenke et al., 1998). As noted by Halle in his foundational study on the MF formalism(Halle, 2009), this approach can also be contextualized within the MF framework of Halle and Wennerström(Halle and Wennerström, 1981).

330 Author contributions. RK, CK and IC devised the project. CK compiled the theoretical considerations. IC and CK performed the numerical simulations. GK and RK assessed the experimental feasibility. GK designed the pulse sequence. IC and CK illustrated the results. CK wrote the manuscript with contributions from all authors.

Competing interests. No competing interests are declared.

Acknowledgements. This work was supported by the Austrian Science Fund FWF (P28359 and P28937).

335 References

360

- Abergel, D. and Bodenhausen, G.: A simple model for NMR relaxation in the presence of internal motions with dynamical coupling, The Journal of Chemical Physics, 121, 761–768, https://doi.org/10.1063/1.1756867, 2004.
- Abergel, D. and Bodenhausen, G.: Predicting internal protein dynamics from structures using coupled networks of hindered rotators, Journal of Chemical Physics, 123, https://doi.org/10.1063/1.2110028, 2005.
- 340 Barnes, C. A., Shen, Y., Ying, J., Takagi, Y., Torchia, D. A., Sellers, J. R., and Bax, A.: Remarkable Rigidity of the Single α-Helical Domain of Myosin-VI As Revealed by NMR Spectroscopy, Journal of the American Chemical Society, 141, 9004–9017, https://doi.org/10.1021/jacs.9b03116, 2019.
 - Bremi, T. and Brüschweiler, R.: Locally Anisotropic Internal Polypeptide Backbone Dynamics by NMR Relaxation, Journal of the American Chemical Society, 119, 6672–6673, https://doi.org/10.1021/ja9708676, 1997.
- 345 Bussell, R. and Eliezer, D.: Residual Structure and Dynamics in Parkinson's Disease-associated Mutants of α-Synuclein*, Journal of Biological Chemistry, 276, 45 996–46 003, https://doi.org/https://doi.org/10.1074/jbc.M106777200, 2001.
 - Carlomagno, T., Maurer, M., Hennig, M., and Griesinger, C.: Ubiquitin Backbone Motion Studied via NHN-C'Cα Dipolar-Dipolar and C'-C'Cα/NHN CSA-Dipolar Cross-Correlated Relaxation, Journal of the American Chemical Society, 122, 5105–5113, https://doi.org/10.1021/ja993845n, 2000.
- 350 Chang, S.-L. and Tjandra, N.: Temperature dependence of protein backbone motion from carbonyl 13C and amide 15N NMR relaxation, Journal of Magnetic Resonance, 174, 43–53, https://doi.org/https://doi.org/10.1016/j.jmr.2005.01.008, 2005.
 - Cisnetti, F., Loth, K., Pelupessy, P., and Bodenhausen, G.: Determination of Chemical Shift Anisotropy Tensors of Carbonyl Nuclei in Proteins through Cross-Correlated Relaxation in NMR, ChemPhysChem, 5, 807–814, https://doi.org/10.1002/cphc.200301041, 2004.
- 355 Clore, G. M. and Iwahara, J.: Theory, Practice, and Applications of Paramagnetic Relaxation Enhancement for the Characterization of Transient Low-Population States of Biological Macromolecules and Their Complexes, Chemical Reviews, 109, 4108–4139, https://doi.org/10.1021/cr900033p, pMID: 19522502, 2009.
 - Clore, G. M., Szabo, A., Bax, A., Kay, L. E., Driscoll, P. C., and Gronenborn, A. M.: Deviations from the simple two-parameter model-free approach to the interpretation of nitrogen-15 nuclear magnetic relaxation of proteins, Journal of the American Chemical Society, 112, 4989–4991, https://doi.org/10.1021/ja00168a070, 1990.
 - Corey, R. B., Pauling, L. C., and Astbury, W. T.: Fundamental dimensions of polypeptide chains, Proceedings of the Royal Society of London. Series B - Biological Sciences, 141, 10–20, https://doi.org/10.1098/rspb.1953.0011, 1953.
 - Cornilescu, G., Marquardt, J. L., Ottiger, M., and Bax, A.: Validation of Protein Structure from Anisotropic Carbonyl Chemical Shifts in a Dilute Liquid Crystalline Phase, Journal of the American Chemical Society, 120, 6836–6837, https://doi.org/10.1021/ja9812610, 1998.
- 365 Daragan, V. A. and Mayo, K. H.: Using the Model Free Approach to Analyze NMR Relaxation Data in Cases of Anisotropic Molecular Diffusion, The Journal of Physical Chemistry B, 103, 6829–6834, https://doi.org/10.1021/jp9911393, 1999.
 - Dayie, K. T. and Wagner, G.: Carbonyl Carbon Probe of Local Mobility in 13C,15N-Enriched Proteins Using High-Resolution Nuclear Magnetic Resonance, Journal of the American Chemical Society, 119, 7797–7806, https://doi.org/10.1021/ja9633880, 1997.

Deschamps, M.: Cross-Correlated Relaxation with Anisotropic Reorientation and Small Amplitude Local Motions, The Journal of Physical

370 Chemistry A, 106, 2438–2445, https://doi.org/10.1021/jp013407e, 2002.

- Deschamps, M. and Bodenhausen, G.: Anisotropy of Rotational Diffusion, Dipole–Dipole Cross-Correlated NMR Relaxation and Angles between Bond Vectors in Proteins, ChemPhysChem, 2, 539–543, https://doi.org/https://doi.org/10.1002/1439-7641(20010917)2:8/9<539::AID-CPHC539>3.0.CO;2-M, 2001.
- Eastman, P., Swails, J., Chodera, J. D., McGibbon, R. T., Zhao, Y., Beauchamp, K. A., Wang, L.-P., Simmonett, A. C., Harrigan, M. P., Stern,
- 375 C. D., Wiewiora, R. P., Brooks, B. R., and Pande, V. S.: OpenMM 7: Rapid development of high performance algorithms for molecular dynamics, PLOS Computational Biology, 13, 1–17, https://doi.org/10.1371/journal.pcbi.1005659, 2017.
 - Ferrage, F., Pelupessy, P., Cowburn, D., and Bodenhausen, G.: Protein Backbone Dynamics through 13C'-13Cα Cross-Relaxation in NMR Spectroscopy, Journal of the American Chemical Society, 128, 11 072–11 078, https://doi.org/10.1021/ja0600577, pMID: 16925424, 2006.
 - Ghose, R., Huang, K., and Prestegard, J. H.: Measurement of Cross Correlation between Dipolar Coupling and Chemi-
- 380 cal Shift Anisotropy in the Spin Relaxation of13C,15N-Labeled Proteins, Journal of Magnetic Resonance, 135, 487 499, https://doi.org/https://doi.org/10.1006/jmre.1998.1602, 1998.
 - Gopal, S. M., Wingbermühle, S., Schnatwinkel, J., Juber, S., Herrmann, C., and Schäfer, L. V.: Conformational Preferences of an Intrinsically Disordered Protein Domain: A Case Study for Modern Force Fields, The Journal of Physical Chemistry B, 125, 24–35, https://doi.org/10.1021/acs.jpcb.0c08702, pMID: 33382616, 2021.
- 385 Grudziąż, K., Zawadzka-Kazimierczuk, A., and Koźmiński, W.: High-dimensional NMR methods for intrinsically disordered proteins studies, Methods, 148, 81–87, https://doi.org/https://doi.org/10.1016/j.ymeth.2018.04.031, nMR Methods of Characterizing Biomolecular Structural Dynamics and Conformational Ensembles, 2018.
 - Halle, B.: The physical basis of model-free analysis of NMR relaxation data from proteins and complex fluids, Journal of Chemical Physics, 131, 1–224 507, https://doi.org/10.1063/1.3269991, 2009.
- 390 Halle, B. and Wennerström, H.: Interpretation of magnetic resonance data from water nuclei in heterogeneous systems, The Journal of Chemical Physics, 75, 1928–1943, https://doi.org/10.1063/1.442218, 1981.
 - Hsu, A., Ferrage, F., and Palmer, A. G.: Analysis of NMR Spin-Relaxation Data Using an Inverse Gaussian Distribution Function, Biophysical Journal, 115, 2301–2309, https://doi.org/10.1016/j.bpj.2018.10.030, 2018.

Idiyatullin, D., Daragan, V. A., and Mayo, K. H.: A New Approach to Visualizing Spectral Density Functions and Deriving Motional Cor-

- 395 relation Time Distributions: Applications to an α-Helix-Forming Peptide and to a Well-Folded Protein, Journal of Magnetic Resonance, 152, 132 – 148, https://doi.org/https://doi.org/10.1006/jmre.2001.2372, 2001.
 - Iwahara, J., Schwieters, C. D., and Clore, G. M.: Ensemble Approach for NMR Structure Refinement against 1H Paramagnetic Relaxation Enhancement Data Arising from a Flexible Paramagnetic Group Attached to a Macromolecule, Journal of the American Chemical Society, 126, 5879–5896, https://doi.org/10.1021/ja031580d, pMID: 15125681, 2004.
- Kadeřávek, P., Zapletal, V., Rabatinová, A., Krásný, L., Sklenář, V., and Žídek, L.: Spectral density mapping protocols for analysis of molecular motions in disordered proteins, Journal of Biomolecular NMR, 58, 193–207, https://doi.org/10.1007/s10858-014-9816-4, 2014.
 Kadeřávek, P., Grutsch, S., Salvi, N., Tollinger, M., Žídek, L., Bodenhausen, G., and Ferrage, F.: Cross-correlated relaxation measurements under adiabatic sweeps: determination of local order in proteins, Journal of Biomolecular NMR, 63, 353–365, https://doi.org/10.1007/s10858-015-9994-8, 2015.
- 405 Kauffmann, C., Zawadzka-Kazimierczuk, A., Kontaxis, G., and Konrat, R.: Using Cross-Correlated Spin Relaxation to Characterize Backbone Dihedral Angle Distributions of Flexible Protein Segments, ChemPhysChem, 22, 18–28, https://doi.org/https://doi.org/10.1002/cphc.202000789, 2021.

- Khan, S., Charlier, C., Augustyniak, R., Salvi, N., Déjean, V., Bodenhausen, G., Lequin, O., Pelupessy, P., and Ferrage, F.: Distribution of Pico- and Nanosecond Motions in Disordered Proteins from Nuclear Spin Relaxation, Biophysical Journal, 109, 988–999,
- 410 https://doi.org/https://doi.org/10.1016/j.bpj.2015.06.069, 2015.
 - Křížová, H., Žídek, L., Stone, M. J., Novotny, M. V., and Sklenář, V.: Temperature-dependent spectral density analysis applied to monitoring backbone dynamics of major urinary protein-I complexed with the pheromone 2-sec-butyl-4,5-dihydrothiazole*, Journal of Biomolecular NMR, 28, 369–384, https://doi.org/10.1023/B:JNMR.0000015404.61574.65, 2004.
 - Kroenke, C. D., Loria, J. P., Lee, L. K., Rance, M., and Palmer, A. G.: Longitudinal and Transverse 1H-15N Dipolar/15N Chemical Shift
- 415 Anisotropy Relaxation Interference: Unambiguous Determination of Rotational Diffusion Tensors and Chemical Exchange Effects in Biological Macromolecules, Journal of the American Chemical Society, 120, 7905–7915, https://doi.org/10.1021/ja9808321, 1998.
 - Kümmerer, F., Orioli, S., Harding-Larsen, D., Hoffmann, F., Gavrilov, Y., Teilum, K., and Lindorff-Larsen, K.: Fitting side-chain NMR relaxation data using molecular simulations, bioRxiv, https://doi.org/10.1101/2020.08.18.256024, 2020.
 - Kämpf, K., Izmailov, S. A., Rabdano, S. O., Groves, A. T., Podkorytov, I. S., and Skrynnikov, N. R.: What Drives 15N Spin Re-
- 420 laxation in Disordered Proteins? Combined NMR/MD Study of the H4 Histone Tail, Biophysical Journal, 115, 2348 2367, https://doi.org/https://doi.org/10.1016/j.bpj.2018.11.017, 2018.
 - Lienin, S. F., Bremi, T., Brutscher, B., Brüschweiler, R., and Ernst, R. R.: Anisotropic Intramolecular Backbone Dynamics of Ubiquitin Characterized by NMR Relaxation and MD Computer Simulation, Journal of the American Chemical Society, 120, 9870–9879, https://doi.org/10.1021/ja9810179, 1998.
- 425 Lipari, G. and Szabo, A.: Model-free approach to the interpretation of nuclear magnetic resonance relaxation in macromolecules. 1. Theory and range of validity, Journal of the American Chemical Society, 104, 4546–4559, https://doi.org/10.1021/ja00381a009, 1982.
 - Loth, K., Pelupessy, P., and Bodenhausen, G.: Chemical Shift Anisotropy Tensors of Carbonyl, Nitrogen, and Amide Proton Nuclei in Proteins through Cross-Correlated Relaxation in NMR Spectroscopy, Journal of the American Chemical Society, 127, 6062–6068, https://doi.org/10.1021/ja0428630, pMID: 15839707, 2005.
- 430 Mantsyzov, A. B., Maltsev, A. S., Ying, J., Shen, Y., Hummer, G., and Bax, A.: A maximum entropy approach to the study of residue-specific backbone angle distributions in α-synuclein, an intrinsically disordered protein, Protein Science, 23, 1275–1290, https://doi.org/https://doi.org/10.1002/pro.2511, 2014.
 - Mantsyzov, A. B., Shen, Y., Lee, J. H., Hummer, G., and Bax, A.: MERA: a webserver for evaluating backbone torsion angle distributions in dynamic and disordered proteins from NMR data, Journal of Biomolecular NMR, 63, 85–95, https://doi.org/10.1007/s10858-015-9971-2, 2015.
- 435 2015. Marcellini, M., Nguyen, M.-H., Mart
 - Marcellini, M., Nguyen, M.-H., Martin, M., Hologne, M., and Walker, O.: Accurate Prediction of Protein NMR Spin Relaxation by Means of Polarizable Force Fields. Application to Strongly Anisotropic Rotational Diffusion, The Journal of Physical Chemistry B, 124, 5103–5112, https://doi.org/10.1021/acs.jpcb.0c01922, pMID: 32501695, 2020.
 - Markwick, P. R. L., Sprangers, R., and Sattler, M.: Local Structure and Anisotropic Backbone Dynamics from Cross-Correlated NMR
- Relaxation in Proteins, Angewandte Chemie International Edition, 44, 3232–3237, https://doi.org/https://doi.org/10.1002/anie.200462495, 2005.
 - Meirovitch, E., Shapiro, Y. E., Polimeno, A., and Freed, J. H.: Protein Dynamics from NMR: The Slowly Relaxing Local Structure Analysis Compared with Model-Free Analysis, The Journal of Physical Chemistry A, 110, 8366–8396, https://doi.org/10.1021/jp056975t, pMID: 16821820, 2006.

- 445 Modig, K. and Poulsen, F. M.: Model-independent interpretation of NMR relaxation data for unfolded proteins: the acid-denatured state of ACBP, Journal of Biomolecular NMR, 42, 163–177, https://doi.org/10.1007/s10858-008-9280-0, 2008.
 - Nodet, G., Abergel, D., and Bodenhausen, G.: Predicting NMR Relaxation Rates in Anisotropically Tumbling Proteins through Networks of Coupled Rotators, ChemPhysChem, 9, 625–633, https://doi.org/10.1002/cphc.200700732, 2008.
- Ottiger, M. and Bax, A.: Determination of Relative N-HN, N-C⁺, C α -C⁺, and C α -H α Effective Bond Lengths in a Protein by NMR in a Dilute Liquid Crystalline Phase, Journal of the American Chemical Society, 120, 12 334–12 341, https://doi.org/10.1021/ja9826791, 1998.
 - Pelupessy, P., Espallargas, G. M., and Bodenhausen, G.: Symmetrical reconversion: measuring cross-correlation rates with enhanced accuracy, Journal of Magnetic Resonance, 161, 258 264, https://doi.org/https://doi.org/10.1016/S1090-7807(02)00190-8, 2003.
 - Pelupessy, P., Ferrage, F., and Bodenhausen, G.: Accurate Measurement of Longitudinal Cross-Relaxation Rates in Nuclear Magnetic Resonance, Journal of Chemical Physics, 126, 134 508, 1–10, https://doi.org/10.1063/1.2715583, 2007.
- 455 Piana, S., Donchev, A. G., Robustelli, P., and Shaw, D. E.: Water Dispersion Interactions Strongly Influence Simulated Structural Properties of Disordered Protein States, The Journal of Physical Chemistry B, 119, 5113–5123, https://doi.org/10.1021/jp508971m, pMID: 25764013, 2015.
 - Piana, S., Robustelli, P., Tan, D., Chen, S., and Shaw, D. E.: Development of a Force Field for the Simulation of Single-Chain Proteins and Protein–Protein Complexes, Journal of Chemical Theory and Computation, 16, 2494–2507, https://doi.org/10.1021/acs.jctc.9b00251,
- 460 pMID: 31914313, 2020.

470

- Rauscher, S., Gapsys, V., Gajda, M. J., Zweckstetter, M., de Groot, B. L., and Grubmüller, H.: Structural Ensembles of Intrinsically Disordered Proteins Depend Strongly on Force Field: A Comparison to Experiment, Journal of Chemical Theory and Computation, 11, 5513–5524, https://doi.org/10.1021/acs.jctc.5b00736, pMID: 26574339, 2015.
- Richert, R. and Blumen, A., eds.: Disorder Effects on Relaxational Processes, Springer Berlin Heidelberg, https://doi.org/10.1007/978-3-642-78576-4, 1994.
 - Robustelli, P., Piana, S., and Shaw, D. E.: Developing a molecular dynamics force field for both folded and disordered protein states, Proceedings of the National Academy of Sciences, 115, E4758–E4766, https://doi.org/10.1073/pnas.1800690115, 2018.
 - Salomon-Ferrer, R., Götz, A. W., Poole, D., Le Grand, S., and Walker, R. C.: Routine Microsecond Molecular Dynamics Simulations with AMBER on GPUs. 2. Explicit Solvent Particle Mesh Ewald, Journal of Chemical Theory and Computation, 9, 3878–3888, https://doi.org/10.1021/ct400314y, pMID: 26592383, 2013.
 - Salvi, N., Abyzov, A., and Blackledge, M.: Multi-Timescale Dynamics in Intrinsically Disordered Proteins from NMR Relaxation and Molecular Simulation, The Journal of Physical Chemistry Letters, 7, 2483–2489, https://doi.org/10.1021/acs.jpclett.6b00885, pMID: 27300592, 2016.
- Salvi, N., Abyzov, A., and Blackledge, M.: Analytical Description of NMR Relaxation Highlights Correlated Dynamics in Intrinsically Dis-
- ordered Proteins, Angewandte Chemie International Edition, 56, 14020–14024, https://doi.org/https://doi.org/10.1002/anie.201706740, 2017.
 - Schanda, P.: Relaxing with liquids and solids A perspective on biomolecular dynamics, Journal of Magnetic Resonance, 306, 180 186, https://doi.org/https://doi.org/10.1016/j.jmr.2019.07.025, 2019.
 - Schwalbe, H., Carlomagno, T., Hennig, M., Junker, J., Reif, B., Richter, C., and Griesinger, C.: [2] Cross-Correlated Relax-
- 480 ation for Measurement of Angles between Tensorial Interactions, in: Nuclear Magnetic Resonance of Biological Macromolecules Part A, edited by James, T. L., Dötsch, V., and Schmitz, U., vol. 338 of *Methods in Enzymology*, pp. 35–81, Academic Press, https://doi.org/https://doi.org/10.1016/S0076-6879(02)38215-6, 2002.

- Shea, J.-E., Best, R. B., and Mittal, J.: Physics-based computational and theoretical approaches to intrinsically disordered proteins, Current Opinion in Structural Biology, 67, 219–225, https://doi.org/10.1016/j.sbi.2020.12.012, 2021.
- 485 Smith, A. A., Ernst, M., and Meier, B. H.: Because the Light is Better Here: Correlation-Time Analysis by NMR Spectroscopy, Angewandte Chemie International Edition, 56, 13 590–13 595, https://doi.org/10.1002/anie.201707316, 2017.
 - Smith, A. A., Ernst, M., Meier, B. H., and Ferrage, F.: Reducing bias in the analysis of solution-state NMR data with dynamics detectors, The Journal of Chemical Physics, 151, 034 102, https://doi.org/10.1063/1.5111081, 2019.
- Stone, J. E., Phillips, J. C., Freddolino, P. L., Hardy, D. J., Trabuco, L. G., and Schulten, K.: Accelerating molecular modeling applications
 with graphics processors, Journal of Computational Chemistry, 28, 2618–2640, https://doi.org/https://doi.org/10.1002/icc.20829, 2007.
 - Stone, J. E., Hardy, D. J., Ufimtsev, I. S., and Schulten, K.: GPU-accelerated molecular modeling coming of age, Journal of Molecular Graphics and Modelling, 29, 116–125, https://doi.org/https://doi.org/10.1016/j.jmgm.2010.06.010, 2010.
 - Tjandra, N., Szabo, A., and Bax, A.: Protein Backbone Dynamics and 15N Chemical Shift Anisotropy from Quantitative Measurement of Relaxation Interference Effects, Journal of the American Chemical Society, 118, 6986–6991, https://doi.org/10.1021/ja960510m, 1996.
- 495 Tugarinov, V., Liang, Z., Shapiro, Y. E., Freed, J. H., and Meirovitch, E.: A Structural Mode-Coupling Approach to 15N NMR Relaxation in Proteins, Journal of the American Chemical Society, 123, 3055–3063, https://doi.org/10.1021/ja003803v, pMID: 11457016, 2001.
 - Vögeli, B.: Comprehensive description of NMR cross-correlated relaxation under anisotropic molecular tumbling and correlated local dynamics on all time scales, The Journal of Chemical Physics, 133, 014 501, https://doi.org/10.1063/1.3454734, 2010.
- Vugmeyster, L., Pelupessy, P., Vugmeister, B. E., Abergel, D., and Bodenhausen, G.: Cross-correlated relaxation in NMR
 of macromolecules in the presence of fast and slow internal dynamics, Comptes Rendus Physique, 5, 377 386, https://doi.org/https://doi.org/10.1016/j.crhy.2004.02.004, highly polarized nuclear spin systems and dipolar interactions in NMR, 2004.
 - Vögeli, B.: The nuclear Overhauser effect from a quantitative perspective, Progress in Nuclear Magnetic Resonance Spectroscopy, 78, 1–46, https://doi.org/https://doi.org/10.1016/j.pnmrs.2013.11.001, 2014.

Vögeli, B. and Yao, L.: Correlated Dynamics between Protein HN and HC Bonds Observed by NMR Cross Relaxation, Journal of the
 American Chemical Society, 131, 3668–3678, https://doi.org/10.1021/ja808616v, pMID: 19235934, 2009.

- Wang, T., Weaver, D. S., Cai, S., and Zuiderweg, E. R. P.: Quantifying Lipari–Szabo modelfree parameters from 13CO NMR relaxation experiments, Journal of Biomolecular NMR, 36, 79–102, https://doi.org/10.1007/s10858-006-9047-4, 2006.
 - Woessner, D. E.: Spin Relaxation Processes in a Two-Proton System Undergoing Anisotropic Reorientation, The Journal of Chemical Physics, 36, 1–4, https://doi.org/10.1063/1.1732274, 1962.
- 510 Xue, Y., Podkorytov, I. S., Rao, D. K., Benjamin, N., Sun, H., and Skrynnikov, N. R.: Paramagnetic relaxation enhancements in unfolded proteins: Theory and application to drkN SH3 domain, Protein Science, 18, 1401–1424, https://doi.org/https://doi.org/10.1002/pro.153, 2009.
 - Ying, J., Roche, J., and Bax, A.: Homonuclear decoupling for enhancing resolution and sensitivity in NOE and RDC measurements of peptides and proteins, Journal of Magnetic Resonance, 241, 97 – 102, https://doi.org/https://doi.org/10.1016/j.jmr.2013.11.006, a special
- 515 "JMR Perspectives" issue: Foresights in Biomolecular Solution-State NMR Spectroscopy From Spin Gymnastics to Structure and Dynamics, 2014.
 - Zerbetto, M., Buck, M., Meirovitch, E., and Polimeno, A.: Integrated Computational Approach to the Analysis of NMR Relaxation in Proteins: Application to ps-ns Main Chain 15N-1H and Global Dynamics of the Rho GTPase Binding Domain of Plexin-B1, The Journal of Physical Chemistry B, 115, 376–388, https://doi.org/10.1021/jp108633v, pMID: 21142011, 2011.

520 Zerze, G. H., Zheng, W., Best, R. B., and Mittal, J.: Evolution of All-Atom Protein Force Fields to Improve Local and Global Properties, The Journal of Physical Chemistry Letters, 10, 2227–2234, https://doi.org/10.1021/acs.jpclett.9b00850, 2019.