

Supplement

Solid-state ^{13}C -NMR spectroscopic determination of sidechain mobilities in zirconium-based metal-organic frameworks

Günter Hempel,¹ Ricardo Kurz,¹ Silvia Paasch,² Kay Saalwächter,¹ and Eike Brunner²

¹*Martin-Luther-Universität Halle-Wittenberg, Institut für Physik – NMR,
Betty-Heimann-Str. 7, 06120 Halle, Germany*

²*Technische Universität Dresden, Fakultät für Chemie und Lebensmittelchemie,
Bioanalytische Chemie, 01062 Dresden, Germany*

S1. LINE DECOMPOSITIONS

See Figs. 1, 2, and 3.

S2. DIPSHIFT DIAGRAMS

In the diagrams in this section, at the left-hand side the data are shown together with the fitted model functions. The right-hand sides show plots of the mean-square deviation χ^2 versus damping constant r and residual dipolar coupling D_{res} . In the latter, the cross marks the position of minimum mean-square deviation χ^2_{min} ; the closed line connects all points where $\chi^2 = 2\chi^2_{\text{min}}$. The evaluation was not executed for all spinning frequencies; it was omitted for such frequencies where the line under consideration had a distance of an integer multiple of spinning frequency to another line.




Figure 1. Decomposition of overlapping middle-ring lines of both protonated carbons and those carbons bound to C≡C in PIZOF-10 (right) and in PIZOF-11 (left).)




Figure 2. Decomposition of overlapping lines in the region 120 ... 140 ppm in PIZOF-2.)




Figure 3. Decomposition of overlapping lines in the region 120 ... 140 ppm in PIZOF-10 and in PIZOF-11.)






Figure 4. PIZOF-2, -15°C, CH of side ring, model function 1

Figure 5. PIZOF-10, -15°C, CH_2 , left component, model function 3Figure 6. PIZOF-10, -15°C, CH_2 , middle component, model function 3







Figure 7. PIZOF-10, -15°C, CH₂, right component, model function 3

Figure 8. PIZOF-10, -15°C, CH of side ring, model function 1

Figure 9. PIZOF-10, 25°C, CH₂ right component, model function 3




Figure 10. PIZOF-10, 25°C, CH₂ 2nd component from right, model function 3




Figure 11. PIZOF-10, 25°C, CH₂ middle component, model function 3




Figure 12. PIZOF-10, 25°C, CH₂ 2nd component from right, model function 3




Figure 13. PIZOF-10, 25°C, CH of middle ring, model function 1




Figure 14. PIZOF-10, 25°C, CH of middle ring, model function 2




Figure 15. PIZOF-10, 25°C, CH of side ring, model function 1




Figure 16. PIZOF-10, 25°C, CH of side ring, model function 2




Figure 17. PIZOF-11, -15°C, δ -CH₂ left component, model function 3







Figure 18. PIZOF-11, -15°C, ε -CH₂, model function 3

Figure 19. PIZOF-11, 30°C, $\gamma\text{-CH}_2$, model function 3Figure 20. PIZOF-11, 30°C, $\delta\text{-CH}_2$, model function 3Figure 21. PIZOF-11, 30°C, $\epsilon\text{-CH}_2$, model function 3



Figure 22. PIZOF-11, 30°C, ω -CH₂, model function 3

Figure 23. PIZOF-11, 25°C, middle ring, model function 1




Figure 24. PIZOF-11, 25°C, side ring, model function 1