

Cell-free Synthesis of Proteins with Selectively ^{13}C -Labelled Methyl Groups from Inexpensive Precursors

Damian Van Raad¹, Gottfried Otting^{1,2} and Thomas Huber¹

¹Australian National University, Research School of Chemistry, Canberra, ACT 2601, Australia

5 ²ARC Centre of Excellence for Innovations in Peptide & Protein Science

Correspondence to: Gottfried Otting (Gottfried.otting@anu.edu.au) and Thomas Huber (t.huber@anu.edu.au)

Abstract. The novel eCell system maintains the activity of the entire repertoire of metabolic *E. coli* enzymes in cell-free protein synthesis. We show that this can be harnessed to produce proteins with selectively ^{13}C -labelled amino acids from inexpensive ^{13}C -labelled precursors. The system is demonstrated with selective ^{13}C -labelling of methyl groups in the
10 proteins ubiquitin and peptidyl-prolyl *cis-trans* isomerase B. Starting from 3- ^{13}C -pyruvate, ^{13}C -HSQC cross-peaks are obtained devoid of one-bond ^{13}C - ^{13}C scalar couplings. Starting from 2- ^{13}C -methyl-acetolactate, single methyl groups of valine and leucine are labelled. Labelling efficiencies are 70% or higher, and the method allows to produce perdeuterated proteins with protonated methyl groups in residue-selective manner. The system uses the isotope-labelled precursors sparingly and is readily scalable.

15 **1 Introduction**

The NMR resonance assignments of high-molecular weight proteins critically depends on the availability of samples enriched with stable isotopes (Tugarinov et al., 2006). Conventional strategies based on uniformly ^{13}C -enriched proteins usually employ [U- ^{13}C]-glucose as the (de facto) only carbon source in minimal media (Ohki and Kainosho, 2008; Filipp et al., 2009). The ^{13}C -enrichment of proteins enables the sensitive recording of heteronuclear correlation spectra such as 2D ^{13}C -HSQC spectra,
20 which are particularly sensitive for methyl groups. Methyl groups play a privileged role in the NMR analysis of large protein systems in solution, as their signals can be observed for macromolecular complexes as large as 1 MDa (Boswell and Latham, 2018). Methyl-bearing amino acids are abundant not only in the hydrophobic core of globular proteins but also in hydrophobic ligand binding pockets (Otten et al., 2010). Methyl groups thus serve as useful probes for the analysis of protein structure, dynamics and function (Schütz and Sprangers, 2020).

25

Among the amino acids with methyl groups, the spectral regions of the methyl groups of isoleucine, leucine and valine (ILV) overlap in a ^{13}C -HSQC spectrum (Rasia et al., 2012). This poses a problem for large proteins, which not only contain many methyl groups but also feature broad NMR signals (Lange et al., 2012). Furthermore, uniformly ^{13}C -labelled proteins feature ^{13}C - ^{13}C couplings, in particular one-bond ^{13}C - ^{13}C couplings, $^1J_{\text{CC}}$, which lead to broad multiplets in the ^{13}C -dimension of ^{13}C -
30 HSQC spectra. Several strategies have been devised to resolve the methyl cross-peaks of ILV residues. (i) Protein samples can

be produced from amino acid mixtures containing only a single amino acid with isotope enrichment. Suitably labelled amino acids are available commercially but can be expensive (Kainosho and Güntert, 2009; Takeda et al., 2010). In many cases, the most affordable versions of ^{13}C -labelled amino acids are uniformly enriched with ^{13}C , which retains the problem of ^{13}C - ^{13}C couplings. (ii) As a compromise between cost and selectivity, selectively ^{13}C -labelled late-stage precursors such as 2-ketobutyrate or 2-ketoisovalerate can be supplied (Goto et al., 1999; Hajduk et al., 2000; Lazarova et al., 2018), which are key intermediates of the biosynthesis of ILV amino acids (Lundström et al., 2007). These precursors are commercially available in selectively ^{13}C - and ^2H -labelled form to produce proteins with single $^{13}\text{CH}_3$ groups in valine, leucine and the δ_1 position of isoleucine in an otherwise perdeuterated background and have proven extraordinarily useful for NMR investigations of high-molecular weight proteins (Tugarinov and Kay, 2005). Precursors close to the final stages of amino acid biosynthesis present a cost-efficient way for labelling proteins with high selectivity (Kasinath et al., 2013; Schörghuber et al., 2018) and, by virtue of specific chemical synthesis, solve the problem of ^{13}C - ^{13}C couplings. (iii) An elegant extension of methyl labelling is presented by the provision of 2- ^{13}C -methylacetolactate in the growth medium, which achieves stereospecific-selective labelling of single methyl groups of valine and leucine (Gans et al., 2010). This approach relies on the activity of several enzymes in the biosynthesis pathways for leucine and valine and thus requires *in vivo* protein production and, consequently, relatively large quantities of the expensive precursor. (iv) One-bond ^{13}C - ^{13}C couplings in uniformly ^{13}C -labelled proteins can also be removed by NMR techniques. For example, ^{13}C - ^1H correlation spectra can be recorded with homonuclear ^{13}C -decoupling in the ^{13}C -dimension, either by recording as a constant-time experiment (Vuister and Bax, 1992) or band-selective decoupling (Behera et al., 2020). However, constant-time experiments sacrifice sensitivity and band-selective decoupling of methyl carbons cannot decouple the ^{13}C multiplet of leucine methyls as the ^{13}C chemical shifts of their coupling partners are too close.

50

Selective methyl labelling by the use of late-stage precursors has become one of the most important approaches for NMR studies of large proteins, having been successfully applied to protein complexes up to 1 MDa (Sprangers and Kay, 2007). The cost of late-stage precursors, however, can become significant when the assignment of the methyl cross-peaks can only be obtained by site-directed mutagenesis. A case in point is the 468-kDa multimeric aminopeptidase PhTET2, where the assignment of the alanine C^βH_3 and isoleucine $\text{C}^\delta\text{H}_3$ groups alone consumed 3.2 L of media with expensive ^{13}C -labelled precursors (Amero et al., 2011). The present work explored the possibility of using earlier precursors of amino acid biosynthesis to produce proteins with $^{13}\text{CH}_3$ groups free from one-bond ^{13}C - ^{13}C couplings and with the option of a background of perdeuteration.

60 The optimal labelling scheme should be amenable to cell-free protein synthesis (CFPS), which uses isotope labelled compounds sparingly (Torizawa et al., 2004). Unfortunately, the biosynthesis of ^{13}C -labelled amino acids is compromised in *in vitro* protein expression systems (Linser et al., 2014), although a limited degree of metabolism can be restored by re-introducing certain cofactors (Jewett et al., 2008). For example, metabolites from glycolysis can be used for energy generation in CFPS, if cofactors such as NAD^+ and CoA are provided (Kim and Swartz, 2001). Energy generation systems have also been

65 based on phosphoenol pyruvate (PEP), as well as pyruvate, glucose and maltodextrin (Caschera and Noireaux, 2015). In our
hands, these systems proved to be more difficult to establish presumably because of their dependence on the activity of multiple
enzymes from the glycolytic pathway.

70 An alternative CFPS approach to proteins with selectively ^{13}C -labelled ILV residues supplements the reaction with the
enzymes required to convert chemically synthesized precursors to the final amino acid. This has been demonstrated with 2-
ketoisovalerate and 4-methyl-2-oxovalerate, adding purified aminotransferase IlvE to catalyse the last step in the biosynthesis
to valine and leucine, respectively (Lazarova et al., 2018). Conducting the CFPS reaction with an earlier precursor such as
methylacetolactate, however, would require additional enzymes to be active.

75 The recently established eCell system solves the problem of maintaining the activity of enzymes required for energy
regeneration in CFPS (Van Raad and Huber, 2021). Here we show that eCells also conserve the activity of biosynthetic
pathways required for amino acid synthesis from simple precursors. eCells are bacterial cells coated with polymers, where the
cell wall has been lysed (Van Raad and Huber, 2021). The resulting cells can no longer replicate, but they still contain all
biomacromolecules required for protein synthesis, while their porous polymer coat gives low-molecular weight compounds
80 free access to the cytosol. eCells thus are ideal vehicles for CFPS. We hypothesized that eCells preserve the activity of all
enzymes involved in amino acid biosynthesis and therefore allow the production of methyl-labelled amino acids from
inexpensive precursors such as 3- ^{13}C -pyruvate or ^{13}C -glucose. In the following we demonstrate the excellent utility of eCells
to produce proteins with selectively ^{13}C -labelled methyl groups in valine and leucine made from pyruvate, 2-methyl-4-
acetolactate and glucose.

85





Figure 1. Biosynthetic pathways of leucine and valine from isotope-labelled precursors. ¹³C-labelled methyl groups are identified by green balls, and methyl groups at natural isotopic abundance are highlighted by orange balls. (A) Biosynthetic pathway starting from 3-¹³C-labelled pyruvate. (B) Stereoselective biosynthetic pathway starting from (*S*)-2-acetolactate. Abbreviations used: KARI, ketol-acid reductoisomerase; DAD, dihydroxy-acid dehydratase; IMS, 2-isopropylmalate synthase; LEU1, 3-isopropylmalate dehydratase; IMDH, 3-isopropylmalate dehydrogenase; BCAT, branched-chain aminotransferase.

2 Materials and methods

95 2.1 Materials

The polyelectrolytes low-molecular weight chitosan (50,000 – 190,000 Da) and sodium alginate were purchased from Merck. The ethyl ester of 2-¹³C-methyl-4-²H₃-acetolactate (ethyl-2-hydroxy-2-¹³C-methyl-3-oxobutanoate) was purchased from Cambridge Isotope Laboratories (CIL; USA). Perdeuterated amino acids were from CIL and Martek Isotopes (USA). 3-¹³C-pyruvate was from Sigma-Aldrich.

100

2.2 Plasmids

A plasmid was constructed with the pCloDF13 origin of replication, the gene of the *E. coli* peptidyl-prolyl *cis-trans* isomerase PpiB with C-terminal His₆-tag under control of the T7 promoter and a spectinomycin resistance gene, generating the plasmid pCDF PpiB CTH. For ubiquitin expression a plasmid was constructed with pCloDF13 origin of replication, the spectinomycin resistance gene and the gene of ubiquitin under control of the T7 promoter (plasmid pCDF Ubi CTH). A *lac* operator was inserted in front of the T7 promoter to reduce background protein expression prior to induction which reduces the ¹³C labelling efficiency.

2.3 Production of eCells

110 *E. coli* Xjb cells were transformed with either pCDF Ubi CTH or pCDF PpiB CTH and grown in LB medium at 37 °C in baffled flasks with shaking at 180 rpm. Endolysin production was induced at the time of inoculation with a final concentration of 3 mM arabinose. Cells were grown to OD₆₀₀ = 0.6, harvested by centrifugation and washed three times with PBS-E buffer (phosphate-buffered saline with 1 mM EDTA, pH 7.4). For coating with chitosan, the cells were resuspended in 0.25 mg/mL chitosan in PBS-E with vigorous shaking for 20 minutes. The cell pellet was washed with PBS-E pH 6.0 three times to remove
115 excess chitosan and then resuspended in 0.25 mg/mL alginate PBS-E solution and subjected to vigorous shaking for 20 minutes. The cells were then washed 3 times with PBS-E pH 6.0, resuspended in PBS-E pH 7.4 and stored at -80 °C. eCell weights reported for different CFPS reactions refer to the sedimented pellet of encapsulated cells following decanting of the wash buffer.

120 2.4 Production of deuterated eCells

5 g sodium pyruvate was dissolved in 50 mL D₂O and the pH adjusted with 0.1 mM KOD to pH 11. The solution was stirred overnight at 95 °C to exchange the protons of pyruvate for deuterium. 500 mL M9 minimal media was prepared in D₂O with 22 mM KH₂PO₄, 42 mM Na₂HPO₄, 8.6 mM NaCl, 18.6 mM NH₄Cl, 500 µL 1 mg/mL thiamine (vitamin B6), 0.1 mM CaCl₂, 250 µL 1000x metal mixture (50 mM FeCl₃, 10 mM MnCl₂, 10 mM ZnSO₄, 2 mM CoCl₂, 2 mM CuCl₂ and 2 mM NiCl₂), 5
125 mM MgSO₄, 3 mM arabinose and 25 mg/mL spectinomycin. The H-D exchange in pyruvate was confirmed by NMR. The deuterated pyruvate was added to the dry mixture of buffer salts and the final pyruvate-M9 medium made up to 500 mL, adjusted to pH 7.2 and filter sterilised prior to inoculation.

XjB(DE3)* cells that had been transformed with pCDF PpiB CTH were trained for production of perdeuterated proteins in a
130 protocol adapted from that reported by Li and Byrd (2022). 15 mL of an overnight starter culture of pCDF PpiB CTH was diluted with 15 mL of deuterated pyruvate-M9 medium and incubated at 37 °C with shaking at 180 rpm. When the OD₆₀₀ reached 1.0, the cells were again diluted with 30 mL of deuterated pyruvate-M9 medium and incubated a second time. Upon reaching OD₆₀₀ = 1.0, the 60 mL culture was spun down, the cells transferred to a 50 mL culture and growth continued
135 overnight at 37 °C with shaking at 180 rpm. The 50 mL culture was added to 400 mL of deuterated pyruvate-M9 medium and left to grow until OD₆₀₀ = 0.75 was reached, after which the cells were encapsulated as described in Section 2.3.

2.5 CFPS systems

The protocol for pyruvate-based CFPS was adapted from the phosphate recycling system by Jewett and Swartz (2004). The CFPS buffer contained 0.9 mM UTP and CTP, 50 mM HEPES, 1.5 mM GTP, 1.5 mM ATP, 0.68 µM folinic acid, 0.64 mM
140 cAMP, 1.7 mM DTT, 3.5 mM of each amino acid (apart from the amino acid(s) to be synthesized by the eCells for isotope enrichment), 60 mM K-Glu, 8 mM Mg-Glu, 2% v/v PEG-8000, 4 mM sodium oxalate, 0.25 mM CoA and 0.33 mM NAD⁺. Roche cOmplete™ Mini protease inhibitor cocktail was added to the CFPS buffer. Of the volume following dissolution of one tablet in 10 mL water, 10% were added to the CFPS reaction. The reaction was conducted with 33 mM pyruvate.

145 The protocol for glucose-based CFPS was likewise adapted from the previously published phosphate recycling system (Jewett and Swartz, 2004). The glucose CFPS buffer contained the same components as the pyruvate-based CFPS protocol, but with 10 mM sodium phosphate dibasic pH 7.5 and without sodium oxalate and pyruvate. The reaction was conducted with 30 mM glucose.

150 The CFPS system using creatine phosphate and creatine kinase as energy source contained the same components as the pyruvate-based CFPS protocol, but without sodium oxalate, pyruvate, CoA or NAD⁺ and adding 250 µg/mL creatine kinase, 80 mM creatine phosphate and 6 mM Mg-Glu instead of 8 mM Mg-Glu.

Following addition of the isotopically enriched precursor, the CFPS buffers for each of these reactions were adjusted to pH 7.5. Frozen aliquots of encapsulated cells were thawed, and the pellet resuspended in CFPS buffer. Lysis of the cell wall occurs spontaneously during thawing (Van Raad and Huber, 2021). CFPS for each experiment was conducted at 37 °C overnight with shaking at 180 rpm.

2.6 Acetolactate labelling

2-¹³C-methyl-4-²H₃-acetolactate as the source for prochiral methyl groups was set free from the ethyl ester by incubating in H₂O with 0.1 M NaOH (NaOD for deuteration experiment) (pH 13) at 37 °C for 30 minutes. The compound was tested both in pyruvate-based CFPS and in CFPS with the creatine-phosphate/creatine kinase system. The CFPS reaction was conducted in 15 mL buffer with 0.1 mM NADP⁺, 3.5 mM 2-¹³C-methyl-4-²H₃-acetolactate and 0.2 mM penoxsulam to inhibit the acetolactate synthase (ALS) enzyme. Ubiquitin was produced from 300 mg eCells and purified using His-Gravitrapp columns (GE Healthcare, USA).

For perdeuterated CFPS, all buffer stocks were dissolved in D₂O, and the pH was adjusted with KOD to pH 7.2. The creatine phosphate based CFPS reaction was conducted in 20 mL D₂O buffer with 5 mM 2-¹³C-methyl-4-²H₃-acetolactate, 0.1 mM NADP⁺, 1 mM of all amino acids in perdeuterated form excluding valine and 0.2 mM penoxsulam to inhibit the acetolactate synthase (ALS) enzyme. PpiB was produced from 800 mg eCells and purified using His-Gravitrapp columns (GE Healthcare, USA).

2.7 Labelling with 3-¹³C-pyruvate or 1-¹³C glucose

Dry 3-¹³C-pyruvate was added to 15 mL CFPS buffer at 33 mM final concentration. Leucine, valine and isoleucine were omitted from the amino acid mixture to allow for ¹³C-labelling of their methyl groups. Ubiquitin and PpiB were expressed using 300 mg eCells and purified using His-Gravitrapp columns. To illustrate the scalability of the reaction, ubiquitin samples were also produced with specific labelling of alanine and valine in 5 mL CFPS buffer using 300 mg eCells with the amino acid of interest omitted from the amino acid mixture. PpiB with ¹³C-labelled valine was produced in 20 mL pyruvate-based CFPS buffer with 1 g eCells with valine omitted from the amino acid mixture.

To test the performance of 1-¹³C-glucose as ¹³C source, dry 1-¹³C glucose was added to 5 mL glucose-based CFPS buffer at 30 mM final concentration. Leucine and valine were omitted from the amino acid mixture to allow for labelling of their methyl groups. To assess the potential of glutamate in the buffer in diluting the ¹³C-label, reactions were conducted with a buffer containing 60 mM K-Glu/8 mM Mg-Glu or 100 mM adipic acid/8 mM MgCl₂.

2.8 NMR spectroscopy and isotope labelling yields


All NMR spectra were recorded at 25 °C using 5 mm NMR tubes and a Bruker 800 or 600 MHz NMR spectrometer equipped with TCI cryoprobes.

190 The isotope labelling efficiency of leucine residues in ubiquitin was assessed by integrating the ^1H -NMR signals of the δ_2 -methyl group of Leu50 and its ^{13}C satellites, which are resolved in the 1D NMR spectrum. For samples without isotope-labelled leucine, the ^{13}C -HSQC cross-peak intensities of the labelled residues were compared with those of an internal standard of 0.1 mM 3- ^{13}C -pyruvate.

195 **3 Results**

3.1 Ubiquitin with ^{13}C -labelled methyl groups in alanine, leucine and valine made from 3- ^{13}C -pyruvate

The biosynthetic methyl labelling strategies were validated using ubiquitin as a model protein. The ^{13}C -label was provided by 3- ^{13}C -pyruvate, which served both as carbon source for amino acid synthesis and energy source for protein production. Omission of leucine and valine from the reaction mixture allows for detection of ^{13}C -labelled valine and leucine produced
200 from pyruvate during the cell-free reaction. As singly ^{13}C -labelled pyruvate contains no neighbouring ^{13}C atoms, the methyl groups of leucines and valines are expected to not show any $^1J_{\text{CC}}$ coupling. This expectation was borne out in the experiment, where the cross-peaks revealed no splittings in the ^{13}C -dimension (Figure 2). Therefore, this labelling scheme delivers better spectral resolution than uniform ^{13}C -labelling schemes, where multiplet splittings due to $^1J_{\text{CC}}$ couplings can be avoided only by specific pulse sequences that compromise sensitivity (Vuister and Bax, 1992; Behera et al., 2022). As the biosynthetic
205 pathways remained intact, the ^{13}C -label was subject to incorporation into a range of amino acids and thus prone to some isotope scrambling. For example, isotopic enrichment was also detected for alanine (due to direct conversion of pyruvate to alanine by alanine-transaminase) and the γ_2 -methyl group of isoleucine. Importantly, however, the labelling efficiency of the isopropyl groups of leucine and valine was high (about 70%).



210 **Figure 2.** ^{13}C -HSQC spectrum of ubiquitin produced from 3- ^{13}C -pyruvate by eCell CFPS, resulting in uniform ^{13}C -labelling of both isopropyl methyl groups of leucine and valine. The protein yield was 0.7 mg from 10 mL eCell CFPS reaction, and the level of isotope labelling was 70%.

The absence of ^{13}C -enrichment of the δ -methyl group of isoleucine is a signature of the biosynthetic pathway, where one
 215 pyruvate molecule is linked with unlabelled acetyl-CoA to form α -ketobutyrate as the precursor of isoleucine, channelling the ^{13}C label into the γ_2 -methyl rather than the δ -methyl group.

As the cell-free reaction was performed in a buffer containing high concentrations of glutamate, we speculated that the degree of isotope labelling could be increased by substituting glutamic acid for adipic acid, which is not easily converted into amino
 220 acids (Jia et al., 2009). Using 100 mM adipic acid/8 mM MgCl_2 instead of 60 mM K-Glu/8 mM Mg-Glu, however, did not increase the labelling efficiency and slightly decreased the protein yield (data not shown).





Figure 3. ^{13}C -HSQC spectra of ubiquitin expressed in a 5 mL CFPS reaction using 300 mg eCells with $3\text{-}^{13}\text{C}$ -pyruvate. (A) Valine was omitted from the amino acid mixture. Protein yield 1.6 mg, ^{13}C -enrichment of the valine methyl groups $>70\%$. (B) Alanine was omitted from the amino acid mixture. Protein yield 1.35 mg, ^{13}C -enrichment of the alanine methyl groups $>85\%$.

Starting from 3-¹³C-pyruvate for biosynthesis, the selectivity of isotope labelling was enhanced by ‘unlabelling’ the amino acids not of interest for labelling, which is achieved simply by adding them to the CFPS reaction at natural isotopic abundance. For example, the ¹³C label was apparent only in the valine methyl groups when only valine was omitted from the amino acid mixture (Figure 3A), and only alanine peaks were observed when only alanine was left out (Figure 3B).

3.2 Ubiquitin with ¹³C-labelled methyl groups in leucine and valine made from 2-¹³C-methyl-acetolactate

2-¹³C-methyl-acetolactate has been shown to allow the *in vivo* production of proteins with stereospecifically labelled isopropyl groups of valine and leucine (Gans et al., 2010). To test the performance of this approach with eCells, a sample of ubiquitin was prepared with the provision of 2-¹³C-methyl-acetolactate and penoxsulam, which is a bactericidal acetolactate synthase (ALS) inhibitor that blocks the biosynthetic conversion of pyruvate to acetolactate, thus abolishing the synthesis of leucine and valine from pyruvate. Both the unlabelled pyruvate and creatine phosphate ATP regeneration systems were used. Both resulted in stereoselective labelling with similar labelling efficiency, highlighting the absence of any significant isotopic dilution by the addition of pyruvate at natural isotopic abundance. Figure 5 shows that the prochiral *S*-methyl groups of ubiquitin were stereoselectively labelled as expected. Although the ALS inhibitor did not entirely prevent the incorporation of unlabelled valine and leucine, presumably due to the unlabelled amino acids already present in the eCells prior to protein production, the isotope labelling efficiency nevertheless reached 70%. Importantly, the eCell system enabled production of this selectively ¹³C-labelled sample from less than 6 mg methyl-acetolactate precursor, and no ¹³C labelling of pro-*R* methyl groups was detectable. The effectiveness of the ALS inhibitor in preventing the production of unlabelled valine and leucine was confirmed by comparison with the isotope labelling efficiency when the CFPS was performed using the widely used ATP-regeneration system with creatine phosphate and creatine kinase (Kigawa et al., 1999; Apponyi et al., 2008). The same isotope labelling efficiency was obtained and the same protein yield (0.7 mg).



250 **Figure 4.** ^{13}C -HSQC spectrum of ubiquitin with labelling of the pro-*S* methyl groups in leucine and valine by using site-specifically ^{13}C -labelled acetolactate in eCell CFPS. Protein yield 0.7 mg, isotope labelling efficiency 70%.

3.3 PpiB with stereospecific ^{13}C -labelled methyl groups in valine

To illustrate the broad applicability of the eCell approach to produce perdeuterated proteins, it was also applied to the *E. coli* peptidyl-prolyl *cis-trans* isomerase B (PpiB), which is a 19 kDa protein. Figure 5A shows the ^{13}C -HSQC cross-peaks of PpiB prepared with 3- ^{13}C -pyruvate while omitting valine. Although the methyl groups of alanine residues are also observed, no two cross-peaks overlap to the extent that they cannot be recognized as separate cross-peaks.

260 Figure 5B shows the ^{13}C -HSQC cross-peaks of perdeuterated PpiB made by eCell CFPS using perdeuterated eCells and 2- ^{13}C -methyl-4- $^2\text{H}_3$ -acetolactate. All amino acids were provided in perdeuterated form and valine was omitted. This resulted in stereoselective labelling of the pro-*S* groups of valine residues in PpiB with a high labelling efficiency (ca. 90%) and adequate yield (1.32 mg). The deuteration level of the protein was high, as shown by a 1D ^1H -NMR spectrum (Figure in SI).





Figure 5. Selective ^{13}C -labelling of the methyl groups of alanine and valine residues in PpiB produced by eCell CFPS. (A) ^{13}C -HSQC spectrum of PpiB produced from $3\text{-}^{13}\text{C}$ -pyruvate with valine omitted. Published assignments are shown (BMRB file 11451). The spectrum

265 also displays the cross-peaks of the γ_1 -methyl groups, but their assignments have not been reported. Protein yield 2.2 mg, isotope labelling efficiency >75%. (B) ^{13}C -HSQC spectrum of PpiB produced from 2- ^{13}C -methyl-acetolactate by CFPS with valine omitted in an eCell CFPS reaction in D_2O using deuterated eCells. The ^{13}C -HSQC spectrum illustrates the selective labelling of the pro-*S*-methyl groups of valine in a perdeuterated protein. The protein yield was 1.3 mg, and the ^{13}C -labelling level was 90%.

270

3.4 eCell CFPS for stereospecific assignments by biosynthetically directed fractional ^{13}C -labelling

Biosynthetic fractional ^{13}C -labelling is a well-established approach to obtain stereospecific assignments of isopropyl methyl groups (Senn et al., 1989; Neri et al., 1989; Schubert et al., 2006). Starting from a mixture of 10% uniformly ^{13}C -labelled glucose and 90% glucose at natural isotopic abundance, the ^{13}C -NMR spectrum of pro-*R* methyl groups displays splittings due to $^1J_{\text{CC}}$ couplings while the pro-*S* methyl groups do not. The approach is inexpensive as only little isotope-labelled glucose is needed. To explore whether eCells maintain the required biosynthetic pathway, a sample of ubiquitin was prepared from a mixture of ^{13}C -labelled and unlabelled glucose. The ^{13}C -HSQC spectrum showed the multiplet finestructures expected for the pro-*R* and pro-*S* methyl groups (Figure 6).

280




Figure 6. ^{13}C -HSQC spectrum of ubiquitin produced by eCell CFPS from a mixture of 10% uniformly ^{13}C -labelled glucose and 90% glucose at natural isotopic abundance. Protein yield 3.8 mg, labelling efficiency below 10%.

285 4 Discussion

The present work shows that biosynthetic pathways naturally established in bacterial cells can be exploited to produce selectively ^{13}C -labelled proteins also in a cell-free reaction to deliver protein in yields sufficient for NMR analysis. In contrast to the preparation of cell extracts by mechanical lysis and high-speed centrifugation, the preparation of eCells uses milder conditions and thus stands a greater chance of preserving the activities of the natural complement of biosynthetic enzymes of
 290 the parent live *E. coli* cells. In this way, eCells combine the potential inherent in biosynthetic pathways with the advantages of conventional CFPS, namely the low requirement of amino acids (Torizawa et al., 2004), compatibility with toxic proteins and facile modification of the solution conditions with regard to compounds small enough to enter the eCells. Importantly, eCells can be produced rapidly and easily. Once prepared, they can be stored at $-80\text{ }^{\circ}\text{C}$ for years without loss of activity.

295 The present work explored the activity of biosynthetic enzymes towards valine and leucine in eCells. As anticipated for
preserved biosynthetic pathways, we readily obtained protein samples with $^{13}\text{CH}_3$ -labelled valine and leucine, where the amino
acids were made from inexpensive pyruvate during the CFPS reaction. Starting from 3- ^{13}C -pyruvate, the scheme maximizes
the spectral resolution of the ^{13}C -HSQC cross-peaks of different methyl groups by avoiding multiplet splittings arising from
large $^1J_{\text{CC}}$ coupling constants. As the biosynthetic pathway from pyruvate to valine and leucine appears intact, it was not
300 surprising to observe also facile conversion of 2- ^{13}C -methyl-4-acetolactate to valine.

Furthermore, the eCell system proved capable of converting glucose into alanine, valine and leucine, allowing the
stereospecific distinction of the isopropyl methyl groups by the classical method of biosynthetic fractional ^{13}C -labelling that
uses an inexpensive mixture of uniformly ^{13}C -labelled glucose with an excess of glucose at natural isotopic abundance (Neri
305 et al., 1989). This scheme allows stereospecific assignments at extraordinarily low cost as far as ^{13}C -labelled glucose is
concerned, but the level of isotope labelling associated with this scheme is intrinsically low, and we therefore prefer 2- ^{13}C -
methyl-4-acetolactate for stereospecific assignments, which also minimizes cross-peak overlap by avoiding $^1J_{\text{CC}}$ multiplet
splittings.

310 Stereospecific ^{13}C -labelling with 2- ^{13}C -methyl-4-acetolactate in *in vivo* protein expression (Gans et al., 2010) has become very
popular, and this precursor is available commercially. (We found the deuterated isotopologue 2- ^{13}C -methyl-4- $^2\text{H}_3$ -acetolactate
to be more readily available than the undeuterated analogue, although the selective ^{13}C -labelling strategy would be beneficial
also without deuteration.) Our results show that eCell CFPS requires only small amounts of 2-methyl-acetolactate to produce
proteins for identification of the pro-*S* methyl groups in ^{13}C -HSQC spectra. To use this labelling scheme in combination with
315 perdeuteration, we supplied all other amino acids in perdeuterated form. While this increases the cost of isotope labelled
material, the labelling scheme is still affordable. Table 1 shows the cost for isotope-labelled precursors used in the experiments
of the present work.

320

325

Table 1. Comparison of precursors and their contribution to the cost of eCell CFPS reaction with 300 mg eCells.¹

¹³ C-labelled precursor	Precursor cost (USD)	Cost of precursor for one reaction (USD)	Total protein yield (mg/mL)	Labelling degree	Position labelled
2- ¹³ C-methyl-4- ² H ₃ -acetolactate	\$1722/g	\$14 ²	0.7	90%	V= γ_2 L= δ_2
3- ¹³ C-pyruvate	\$866/g	\$34	0.8	70%	V= γ_2, γ_1 L= δ_2, δ_1 I= γ_2
10% [U- ¹³ C]-glucose + 90% unlabelled glucose	\$258/g	\$2	3.8	10%	V= γ_2^3
1- ¹³ C-glucose	\$282/g	\$14	1.4	44%	V= γ_2, γ_1 L= δ_2, δ_1

¹ Prices from Cambridge Isotope Laboratories (<https://www.isotope.com>), Omicron Biochemicals Inc. (<https://www.omicronbio.com>) and Apollo Scientific (D₂O; <https://store.apolloscientific.co.uk>), accessed 3rd April 2023.

330 ² Additional isotope costs were for perdeuterated amino acids (\$222; Table S1) and the D₂O (\$838/L, \$377 for the 0.45 L cell culture used) for growing perdeuterated *E. coli* cells.

³ eCells can synthesize leucine from glucose (Figure S2). Therefore, stereospecific isotope labelling of the δ_2 position of leucine may also be achieved but this was not tested experimentally.

335

To the best of our knowledge, purified ILV amino acids with stereospecific ¹³C-enrichment of single methyl groups are not commercially available. As an alternative, Linser et al. (2014) showed that CFPS reactions can be conducted with an amino acid mixture produced by hydrolysis of a suitably isotope labelled protein expressed *in vivo*. Also in this approach, however, some amino acids need to be supplied in purified form if they are degraded during hydrolysis of the labelled protein.

340 Assembling the amino acid mixture from commercially available individual components is less laborious and offers the important advantage that a single amino acid can be omitted and thus targeted for production by biosynthesis. In this way we obtained high levels of ¹³C incorporation (90%) and deuteration (estimated to be >95%), which are comparable with *in vivo* protein preparations and favourable for good sensitivity of NMR experiments of large protein complexes (O'Brien et al., 2018). In practice, the economical use of amino acids in the eCell CFPS reaction meant that the cost of D₂O used for producing
345 perdeuterated eCells (\$377) exceeded that of the perdeuterated amino acids added in the CFPS reaction (\$222, Table S1).

Pyruvate plays a central role in bacterial biosynthesis and, as shown in the present work, singly ^{13}C -labelled pyruvate is suitable as a relatively inexpensive precursor for labelling methyl groups of leucine and valine with high levels of ^{13}C -enrichment. If, at the same time, unlabelled leucine or valine is provided in the CFPS reaction to suppress their respective cross-peaks, the cross-peaks of the amino acid omitted can be observed selectively. The increased spectral resolution afforded by this scheme is particularly beneficial for larger proteins. Furthermore, inactivation of transaminases by reduction with NaBH_4 (Su et al., 2011) may allow extending this approach to the selective ^{15}N -labelling of amino acids from ^{15}N -ammonium salt. These experiments are currently in progress.

In principle, using 1- ^{13}C -glucose as the carbon source delivers the same selectivity of isotope labelling as 3- ^{13}C -pyruvate (Lundström et al., 2007) but, as glycolysis breaks the glucose down into 3- ^{13}C -pyruvate and unlabelled pyruvate, glucose simultaneously labelled in the 1 and 6 position is required to avoid the dilution with unlabelled pyruvate (Loquet et al., 2011). We therefore prefer 3- ^{13}C -pyruvate.

As pyruvate can be converted to alanine by a single enzyme, it is difficult to suppress the cross-peaks of the C^βH_3 groups of alanine when starting from ^{13}C -labelled pyruvate. The addition of an excess of unlabelled alanine to the reaction would dilute the labelled pyruvate with unlabelled pyruvate, and inhibition of the alanine aminotransferase by reduction with NaBH_4 would also inhibit the transaminase that installs the amino group on leucine and valine by transfer from glutamate. We therefore propose to identify the alanine cross-peaks with a sample, where the isotope labelling of leucine and valine is suppressed by provision of these amino acids in unlabelled form (Figure 3B).

Starting from pyruvate, we found it difficult to achieve ^{13}C -labelling efficiencies much above 70%. We attribute this to an isotope dilution effect due to a pool of unlabelled amino acids present in the eCells. Attempts to dialyse eCells in a large volume of buffer for an extended period of time reduced the protein yield as the eCells lose activity by gradually leaking biomacromolecules (Van Raad and Huber, 2021). Notably, proteins produced *in vivo* from various ^{13}C -labelled glucose isotopomers are likewise subject to isotopic dilution, and examples with ~45% labelling efficiency have been reported (Lundström et al., 2009; Loquet et al., 2011; Weininger, 2017).

As proteins slowly leak through the porous polymer coating, the lifetime of eCells is limited to about 8 h at 37 °C, which limits protein yields. We note, however, that *in vivo* protein expression from selectively labelled precursors cannot be conducted for too long either, if isotope scrambling by precursor recycling is to be avoided (Kurauskas et al., 2017).

380

5 Conclusions

In summary, the eCell platform opens new possibilities for the selective ^{13}C -enrichment of methyl groups in proteins. It combines high levels of isotope enrichment with low cost of isotope-enriched precursors. The protocol for eCell preparation is uncomplicated, and the ready accessibility of the interior space of eCells to low-molecular-weight compounds provides
385 ready control over the chemical environment, so that different isotope labelling is achieved simply by use of different reaction buffers. In contrast to conventional CFPS based on dialysis systems, where protein yields depend on good contact between inside and outside buffers and, therefore, the geometry of the setup, the eCell system can readily be scaled in volume. We anticipate that it will find many more uses beyond those demonstrated in the present work.

390 **Data availability.** The NMR data are available at <https://doi.org/10.5281/zenodo.7662927> (Van Raad et al., 2023).

Supplement. The supplement contains the nucleotide sequences of genes used in this work, and high-resolution mass spectrum of the deuterated PpiB sample.

395 **Author contributions.** DVR initiated the project and performed all biochemical experiments. GO performed all NMR experiments. GO and TH coordinated the project and contributed advice towards experimental design. All three authors contributed to the final manuscript.

Competing interests. The Australian National University holds a patent related to the production and use of eCells
400 (PCT/AU2020/050050) and share financial return from the patent with the inventors (TH and DVR). At least one of the (co-)authors is a member of the editorial board of *Magnetic Resonance*.

Acknowledgements. Gottfried Otting thanks the Australian Research Council, for a Laureate Fellowship (grant no. FL170100019).
405

Financial support. This research has been supported by the Australian Research Council (grant no. FL170100019, DP200100348, DP21010088) and the Australian Research Council Centre of Excellence for Innovations in Peptide and Protein Science (grant no. CE200100012).

410 **NMR spectra are available on Zenodo:** <https://doi.org/10.5281/zenodo.7662927> (Van Raad et al., 2023)

References

- Amero, C., Asunción Durá, M., Noirclerc-Savoie, M., Perollier, A., Gallet, B., Plevin, M. J., Vernet, T., Franzetti, B., and Boisbouvier, J.: A systematic mutagenesis-driven strategy for site-resolved NMR studies of supramolecular assemblies, *J. Biomol. NMR*, 50, 229–236, <https://doi.org/10.1007/s10858-011-9513-5>, 2011.
- 415 Apponyi, M. A., Ozawa, K., Dixon, N. E., and Otting, G.: Cell-free protein synthesis for analysis by NMR spectroscopy, *Methods Mol. Biol.*, 426, 257–268, https://doi.org/10.1007/978-1-60327-058-8_16, 2008.
- Behera, S. P., Dubey, A., Chen, W. N., De Paula, V. S., Zhang, M., Sgourakis, N. G., Bermel, W., Wagner, G., Coote, P. W., and Arthanari, H.: Nearest-neighbor NMR spectroscopy: categorizing spectral peaks by their adjacent nuclei, *Nat. Commun.*, 11, 5547, <https://doi.org/10.1038/s41467-020-19325-4>, 2020.
- 420 Boswell, Z. K. and Latham, M. P.: Methyl-based NMR spectroscopy methods for uncovering structural dynamics in large proteins and protein complexes, *Biochemistry*, 58, 144–155, <https://doi.org/10.1021/acs.biochem.8b00953>, 2018.
- Caschera, F. and Noireaux, V.: A cost-effective polyphosphate-based metabolism fuels an all *E. coli* cell-free expression system, *Metabol. Engin.*, 27, 29–37, <https://doi.org/10.1016/j.ymben.2014.10.007>, 2015.
- Filipp, F. V., Sinha, N., Jairam, L., Bradley, J., and Opella, S. J.: Labeling strategies for ^{13}C -detected aligned-sample solid-
425 state NMR of proteins, *J. Magn. Reson.*, 201, 121–130, <https://doi.org/10.1016/j.jmr.2009.08.012>, 2009.
- Gans, P., Hamelin, O., Sounier, R., Ayala, I., Durá, M. A., Amero, C. D., Noirclerc-Savoie, M., Franzetti, B., Plevin, M. J., and Boisbouvier, J.: Stereospecific isotopic labeling of methyl groups for NMR spectroscopic studies of high-molecular-weight proteins, *Angew. Chem. Int. Ed.*, 49, 1958–1962, <https://doi.org/10.1002/anie.200905660>, 2010.
- Goto, N. K., Gardner, K. H., Mueller, G. A., Willis, R. C., and Kay, L. E.: A robust and cost-effective method for the production
430 of Val, Leu, Ile (δ^1) methyl-protonated ^{15}N -, ^{13}C -, ^2H -labeled proteins, *J. Biomol. NMR*, 13, 369–374, <https://doi.org/10.1023/a:1008393201236>, 1999.
- Hajduk, P. J., Augeri, D. J., Mack, J., Mendoza, R., Yang, J., Betz, S. F., and Fesik, S. W.: NMR-based screening of proteins containing ^{13}C -labeled methyl groups, *J. Am. Chem. Soc.*, 122, 7898–7904, <https://doi.org/10.1021/ja000350l>, 2000.
- Jewett, M. C. and Swartz, J. R.: Mimicking the *Escherichia coli* cytoplasmic environment activates long-lived and efficient
435 cell-free protein synthesis, *Biotechnol. Bioeng.*, 86, 19–26, <https://doi.org/10.1002/bit.20026>, 2004.
- Jewett, M. C., Calhoun, K. A., Voloshin, A., Wu, J. J., and Swartz, J. R.: An integrated cell-free metabolic platform for protein production and synthetic biology, *Mol. Sys. Biol.*, 4, 220, <https://doi.org/10.1038/msb.2008.57>, 2008.
- Jia, X., Ozawa, K., Loscha, K., and Otting, G.: Glutarate and N-acetyl-L-glutamate buffers for cell-free synthesis of selectively
 ^{15}N -labelled proteins, *J. Biomol. NMR*, 44, 59–67, <https://doi.org/10.1007/s10858-009-9315-1>, 2009.
- 440 Kainosho, M. and Güntert, P.: SAIL – stereo-array isotope labeling, *Quart. Rev. Biophys.*, 42, 247–300, <https://doi.org/10.1017/S0033583510000016>, 2009.
- Kasinath, V., Valentine, K. G., and Wand, A. J.: A ^{13}C labeling strategy reveals a range of aromatic side chain motion in calmodulin, *J. Am. Chem. Soc.*, 135, 9560–9563, <https://doi.org/10.1021/ja4001129>, 2013.

- Kigawa, T., Yabuki, T., Yoshida, Y., Tsutsui, M., Ito, Y., Shibata, T., and Yokoyama, S.: Cell-free production and stable-
445 isotope labeling of milligram quantities of proteins. *FEBS Lett.*, 442, 15–19, [https://doi.org/10.1016/S0014-5793\(98\)01620-2](https://doi.org/10.1016/S0014-5793(98)01620-2), 1999.
- Kim, D. M. and Swartz, J. R.: Regeneration of adenosine triphosphate from glycolytic intermediates for cell-free protein synthesis, *Biotechnol. Bioeng.*, 74, 309–316, <https://doi.org/10.1002/bit.1121>, 2001.
- Kurauskas, V., Schanda, P., and Sounier, R.: Methyl-specific isotope labelling strategies for NMR studies of membrane
450 proteins, *Meth. Mol. Biol.*, 1635, 109–123, https://doi.org/10.1007/978-1-4939-7151-0_6, 2017.
- Lange, O. F., Rossi, P., Sgourakis, N. G., Song, Y., Lee, H. W., Aramini, J. M., Ertekin, A., Xiao, R., Acton, T. B., Montelione, G. T., and Baker, D.: Determination of solution structures of proteins up to 40 kDa using CS-Rosetta with sparse NMR data from deuterated samples, *Proc. Natl. Acad. Sci.*, 109, 10873–10878, <https://doi.org/10.1073/pnas.1203013109>, 2012.
- Lazarova, M., Löhr, F., Rues, R.-B., Kleebach, R., Dötsch, V., and Bernhard, F.: Precursor-based selective methyl labeling of
455 cell-free synthesized proteins, *ACS Chem. Biol.*, 13, 2170–2178, <https://doi.org/10.1021/acscchembio.8b00338>, 2018.
- Li, J. and Byrd, R. A.: A simple protocol for the production of highly deuterated proteins for biophysical studies, *J. Biol. Chem.*, 298, 102253, <https://doi.org/10.1016/j.jbc.2022.102253>, 2022.
- Linser, R., Gelev, V., Hagn, F., Arthanari, H., Hyberts, S. G., and Wagner, G.: Selective methyl labeling of eukaryotic membrane proteins using cell-free expression, *J. Am. Chem. Soc.*, 136, 11308–11310, <https://doi.org/10.1021/ja504791j>,
460 2014.
- Loquet, A., Lv, G., Giller, K., Becker, S., and Lange, A.: ^{13}C spin dilution for simplified and complete solid-state NMR resonance assignment of insoluble biological assemblies, *J. Am. Chem. Soc.*, 133, 4722–4725, <https://doi.org/10.1021/ja200066s>, 2011.
- Lundström, P., Teilum, K., Carstensen, T., Bezsonova, I., Wiesner, S., Hansen, D. F., Religa, T. L., Akke, M., and Kay, L. E.:
465 Fractional ^{13}C enrichment of isolated carbons using $[1-^{13}\text{C}]$ -or $[2-^{13}\text{C}]$ -glucose facilitates the accurate measurement of dynamics at backbone C^α and side-chain methyl positions in proteins, *J. Biomol. NMR*, 38, 199–212, <https://doi.org/10.1007/s10858-007-9158-6>, 2007.
- Lundström, P., Lin, H., and Kay, L. E.: Measuring $^{13}\text{C}^\beta$ chemical shifts of invisible excited states in proteins by relaxation dispersion NMR spectroscopy, *J. Biomol. NMR.*, 44, 139–155, <https://doi.org/10.1007/s10858-009-9321-3>, 2009.
- 470 Neri, D., Szyperski, T., Otting, G., Senn, H., and Wüthrich K.: Stereospecific nuclear magnetic resonance assignments of the methyl groups of valine and leucine in the DNA-binding domain of the 434 repressor by biosynthetically directed fractional ^{13}C labeling, *Biochemistry*, 28, 7510–7516, <https://doi.org/10.1021/bi00445a003>, 1989.
- Ohki, S. Y. and Kainosho, M.: Stable isotope labeling methods for protein NMR spectroscopy, *Prog. Nucl. Magn. Reson. Spectrosc.*, 53, 208–226, <https://doi.org/10.1016/j.pnmrs.2008.01.003>, 2008.
- 475 O'Brien, E. S., Lin, D. W., Fuglestad, B., Stetz, M. A., Gosse, T., Tommos, C., and Wand, A. J. Improving yields of deuterated, methyl labeled protein by growing in H_2O , *J. Biomol. NMR*, 71, 263–273, <https://doi.org/10.1007/s10858-018-0200-7>, 2018.

- Otten, R., Chu, B., Krewulak, K. D., Vogel, H. J., and Mulder, F. A. A.: Comprehensive and cost-effective NMR spectroscopy of methyl groups in large proteins, *J. Am. Chem. Soc.*, 132, 2952–2956, <https://doi.org/10.1021/ja907706a>, 2010.
- 480 Rasia, R. M., Brutscher, B., and Plevin, M. J.: Selective isotopic unlabeled of proteins using metabolic precursors: application to NMR assignment of intrinsically disordered proteins, *ChemBioChem.*, 13, 732–739, <https://doi.org/10.1002/cbic.201100678>, 2012.
- Schörghuber, J., Geist, L., Platzer, G., Feichtinger, M., Bisaccia, M., Scheibelberger, L., Weber, F., Konrat, R., and Lichtenecker, R. J.: Late metabolic precursors for selective aromatic residue labeling, *J. Biomol. NMR.*, 71, 129–140, 485 <https://doi.org/10.1007/s10858-018-0188-z>, 2018.
- Schubert, M., Manolikas, T., Rogowski, M., and Meier, B. H.: Solid-state NMR spectroscopy of 10% ¹³C labeled ubiquitin: spectral simplification and stereospecific assignment of isopropyl groups, *J. Biomol. NMR.*, 35, 167–173, <https://doi.org/10.1007/s10858-006-9025-x>, 2006.
- Schütz, S. and Sprangers, R.: Methyl TROSY spectroscopy: A versatile NMR approach to study challenging biological 490 systems. *Prog. Nucl. Magn. Reson. Spectrosc.*, 116, 56–84, <https://doi.org/10.1016/j.pnmrs.2019.09.004>, 2020.
- Senn, H., Werner, B., Messerle, B. A., Weber, C., Traber, R., and Wüthrich, K. Stereospecific assignment of the methyl ¹H NMR lines of valine and leucine in polypeptides by nonrandom ¹³C labelling, *FEBS Lett.*, 249, 113–118, [https://doi.org/10.1016/0014-5793\(89\)80027-4](https://doi.org/10.1016/0014-5793(89)80027-4), 1989.
- Sprangers, R. and Kay, L. E.: Quantitative dynamics and binding studies of the 20S proteasome by NMR, *Nature*, 445, 618– 495 622, <https://doi.org/10.1038/nature05512>, 2007.
- Su, X. C., Loh, C. T., Qi, R., and Otting, G.: Suppression of isotope scrambling in cell-free protein synthesis by broadband inhibition of PLP enzymes for selective ¹⁵N-labelling and production of perdeuterated proteins in H₂O, *J. Biomol. NMR*, 50, 35–42, <https://doi.org/10.1007/s10858-011-9477-5>, 2011.
- Takeda, M., Ono A. M., Terauchi, T., and Kainosho, M.: Application of SAIL phenylalanine and tyrosine with alternative 500 isotope-labeling patterns for protein structure determination, *J. Biomol. NMR*, 46, 45–49, <https://doi.org/10.1007/s10858-009-9360-9>, 2010.
- Torizawa, T., Shimizu, M., Taoka, M., Miyano, H., and Kainosho, M.: Efficient production of isotopically labeled proteins by cell-free synthesis: a practical protocol, *J. Biomol. NMR*, 30, 311–325, <https://doi.org/10.1007/s10858-004-3534-2>, 2004.
- Tugarinov, V., Kanelis, V., and Kay, L. E.: Isotope labeling strategies for the study of high-molecular-weight proteins by 505 solution NMR spectroscopy, *Nat. Protoc.*, 1, 749–754, <https://doi.org/10.1038/nprot.2006.101>, 2006.
- Tugarinov, V. and Kay, L. E.: Methyl groups as probes of structure and dynamics in NMR studies of high-molecular-weight proteins. *Chembiochem.*, 6, 1567–1577, <https://doi.org/10.1002/cbic.200500110>, 2005.
- Van Raad, D. and Huber, T.: *In vitro* protein synthesis in semipermeable artificial cells, *ACS Synth. Biol.*, 10, 1237–1244, <https://doi.org/10.1021/acssynbio.1c00044>, 2021.

- 510 Van Raad, D., Otting, G., and Huber, T. NMR Spectra relating to the publication, Version 1, Zenodo [Cell-free Synthesis of Proteins with Selectively ^{13}C -Labelled Methyl Groups from Inexpensive Precursors-Dataset], <https://doi.org/10.5281/zenodo.7662927>, 2023.
- Vuister, G. W. and Bax, A.: Resolution enhancement and spectral editing of uniformly ^{13}C -enriched proteins by homonuclear broadband ^{13}C decoupling, *J. Magn. Reson.*, 98, 428–435, [https://doi.org/10.1016/0022-2364\(92\)90144-V](https://doi.org/10.1016/0022-2364(92)90144-V), 1992.
- 515 Weininger, U.: Site-selective ^{13}C labeling of proteins using erythrose, *J. Biomol. NMR*, 67, 191–200, <https://doi.org/10.1007/s10858-017-0096-7>, 2017.